

Anti-Histamine H3 Receptor Antibody

Rabbit polyclonal antibody to Histamine H3 Receptor Catalog # AP60094

Specification

Anti-Histamine H3 Receptor Antibody - Product Information

Application WB, IHC
Primary Accession Q9Y5N1
Other Accession P58406

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 48671

Anti-Histamine H3 Receptor Antibody - Additional Information

Gene ID 11255

Other Names

GPCR97; Histamine H3 receptor; H3R; HH3R; G-protein coupled receptor 97

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Histamine H3 Receptor. The exact sequence is proprietary.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200) IHC~~1:100~500

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-Histamine H3 Receptor Antibody - Protein Information

Name HRH3

Synonyms GPCR97

Function

The H3 subclass of histamine receptors could mediate the histamine signals in CNS and peripheral nervous system. Signals through the inhibition of adenylate cyclase and displays high constitutive activity (spontaneous activity in the absence of agonist). Agonist stimulation of isoform 3 neither modified adenylate cyclase activity nor induced intracellular calcium mobilization.



Cellular Location

Cell membrane; Multi-pass membrane protein.

Tissue Location


Expressed predominantly in the CNS, with the greatest expression in the thalamus and caudate nucleus. The various isoforms are mainly coexpressed in brain, but their relative expression level varies in a region-specific manner. Isoform 3 and isoform 7 are highly expressed in the thalamus, caudate nucleus and cerebellum while isoform 5 and isoform 6 show a poor expression. Isoform 5 and isoform 6 show a high expression in the amygdala, substantia nigra, cerebral cortex and hypothalamus. Isoform 7 is not found in hypothalamus or substantia nigra

Anti-Histamine H3 Receptor Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Histamine H3 Receptor Antibody - Images



Western blot analysis of Histamine H3 Receptor expression in HEK293T (A), Hela (B), H460 (C) whole cell lysates.





Immunohistochemical analysis of Histamine H3 Receptor staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-Histamine H3 Receptor Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Histamine H3 Receptor. The exact sequence is proprietary.