

Anti-USP32 Antibody
Rabbit polyclonal antibody to USP32
Catalog # AP60122

Specification

Anti-USP32 Antibody - Product Information

Application	WB
Primary Accession	Q8NFA0
Reactivity	Human, Mouse, Rat, SARS, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	181656

Anti-USP32 Antibody - Additional Information

Gene ID 84669

Other Names

USP10; Ubiquitin carboxyl-terminal hydrolase 32; Deubiquitinating enzyme 32; Renal carcinoma antigen NY-REN-60; Ubiquitin thioesterase 32; Ubiquitin-specific-processing protease 32

Target/Specificity

Recognizes endogenous levels of USP32 protein.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-USP32 Antibody - Protein Information

Name USP32

Synonyms USP10

Function

Deubiquitinase that can remove conjugated ubiquitin from target proteins, such as RAB7A and LAMTOR1 (PubMed: <http://www.uniprot.org/citations/36476874> target="_blank">36476874). Acts as a positive regulator of the mTORC1 signaling by mediating deubiquitination of LAMTOR1, thereby promoting the association between LAMTOR1 and the lysosomal V-ATPase complex and subsequent activation of the mTORC1 complex (PubMed: <http://www.uniprot.org/citations/36476874> target="_blank">36476874).

Cellular Location


Golgi apparatus membrane; Lipid-anchor

Anti-USP32 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-USP32 Antibody - Images



Western blot analysis of USP32 expression in A549 (A), MCF7 (B), PC12 (C), AML12 (D) whole cell lysates.



Immunohistochemical analysis of USP32 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-USP32 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human USP32. The exact sequence is proprietary.