

**Anti-CABLES1 Antibody**  
**Rabbit polyclonal antibody to CABLES1**  
**Catalog # AP60124****Specification**

---

**Anti-CABLES1 Antibody - Product Information**

Application	WB, IF/IC, IHC
Primary Accession	<a href="#">Q8TDN4</a>
Other Accession	<a href="#">Q9ESJ1</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67599

**Anti-CABLES1 Antibody - Additional Information****Gene ID** 91768**Other Names**

CABLES; CDK5 and ABL1 enzyme substrate 1; Interactor with CDK3 1; Ik3-1

**Target/Specificity**

Recognizes endogenous levels of CABLES1 protein.

**Dilution**

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500)

IF/IC~~N/A

IHC~~1:100~500

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-CABLES1 Antibody - Protein Information****Name** CABLES1**Synonyms** CABLES**Function**

Cyclin-dependent kinase binding protein. Enhances cyclin- dependent kinase tyrosine phosphorylation by nonreceptor tyrosine kinases, such as that of CDK5 by activated ABL1, which leads to increased CDK5 activity and is critical for neuronal development, and that of CDK2 by WEE1, which leads to decreased CDK2 activity and growth inhibition. Positively affects neuronal outgrowth. Plays a role as a regulator for p53/p73-induced cell death (By similarity).

**Cellular Location**

Nucleus. Cytoplasm. Note=Located in the cell body and proximal region of the developing axonal shaft of immature neurons. Located in axonal growth cone, but not in the distal part of the axon shaft or in dendritic growth cone of mature neurons (By similarity).


**Tissue Location**

Expressed in breast, pancreas, colon, head and neck (at protein level). Strongly decreased in more than half of cases of atypical endometrial hyperplasia and in more than 90% of endometrial cancers.


**Anti-CABLES1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-CABLES1 Antibody - Images**

Western blot analysis of CABLES1 expression in Hela (A), H446 (B) whole cell lysates.



Immunohistochemical analysis of CABLES1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of CABLES1 staining in SHSY5Y cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

#### **Anti-CABLES1 Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human CABLES1. The exact sequence is proprietary.