

## Anti-HSD17B4 Antibody

Rabbit polyclonal antibody to HSD17B4 Catalog # AP60180

### Specification

## Anti-HSD17B4 Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Calculated MW WB <u>P51659</u> <u>P51660</u> Human, Mouse, Rat, Monkey Rabbit Polyclonal 79686

## **Anti-HSD17B4 Antibody - Additional Information**

Gene ID 3295

**Other Names** EDH17B4; Peroxisomal multifunctional enzyme type 2; MFE-2; 17-beta-hydroxysteroid dehydrogenase 4; 17-beta-HSD 4; D-bifunctional protein; DBP; Multifunctional protein 2; MPF-2

**Target/Specificity** KLH-conjugated synthetic peptide encompassing a sequence within the center region of human HSD17B4. The exact sequence is proprietary.

Dilution WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

## Anti-HSD17B4 Antibody - Protein Information

Name HSD17B4 (<u>HGNC:5213</u>)

Synonyms EDH17B4, SDR8C1

#### Function

Bifunctional enzyme acting on the peroxisomal fatty acid beta-oxidation pathway. Catalyzes two of the four reactions in fatty acid degradation: hydration of 2-enoyl-CoA (trans-2-enoyl-CoA) to produce (3R)-3-hydroxyacyl-CoA, and dehydrogenation of (3R)-3- hydroxyacyl-CoA to produce 3-ketoacyl-CoA (3-oxoacyl-CoA), which is further metabolized by SCPx. Can use straight-chain and branched-chain fatty acids, as well as bile acid intermediates as substrates.



**Cellular Location** Peroxisome.

#### **Tissue Location**


Present in many tissues with highest concentrations in liver, heart, prostate and testis

## Anti-HSD17B4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

## Anti-HSD17B4 Antibody - Images



Western blot analysis of HSD17B4 expression in HEK293T (A), H1688 (B), H1792 (C), mouse liver (D), rat liver (E) whole cell lysates.

# Anti-HSD17B4 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human HSD17B4. The exact sequence is proprietary.