

Anti-GLURD2 Antibody
Rabbit polyclonal antibody to GLURD2
Catalog # AP60300**Specification**

Anti-GLURD2 Antibody - Product Information

Application	WB, IF
Primary Accession	O43424
Other Accession	Q61625
Reactivity	Human, Mouse, Rat, Zebrafish
Host	Rabbit
Clonality	Polyclonal
Calculated MW	113356

Anti-GLURD2 Antibody - Additional Information**Gene ID** 2895**Other Names**

GLURD2; Glutamate receptor ionotropic, delta-2; GluD2; GluR delta-2 subunit

Target/Specificity

Recognizes endogenous levels of GLURD2 protein.

Dilution

WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500)

IF~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-GLURD2 Antibody - Protein Information**Name** GRID2**Synonyms** GLURD2**Function**

Receptor for glutamate. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. The postsynaptic actions of Glu are mediated by a variety of receptors that are named according to their selective agonists. Promotes synaptogenesis and mediates the D-Serine-dependent long term depression signals and AMPA receptor endocytosis of cerebellar parallel fiber- Purkinje cell (PF-PC) synapses through the beta-NRX1-CBLN1-GRID2 triad complex (PubMed:27418511).


Cellular Location

Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane; Multi-pass membrane protein


Anti-GLURD2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-GLURD2 Antibody - Images

Western blot analysis of GLURD2 expression in zebrafish (A) whole cell lysates.



Immunofluorescent analysis of GLURD2 staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

DAPI was used to stain the cell nuclei (blue).

Anti-GLURD2 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human GLURD2. The exact sequence is proprietary.