

**Anti-Nogo Antibody**  
**Rabbit polyclonal antibody to Nogo**  
**Catalog # AP60392****Specification**

---

**Anti-Nogo Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">Q9NQC3</a>
Other Accession	<a href="#">Q99P72</a>
Reactivity	Human, Mouse, Rat, Rabbit, Monkey, Bovine, SARS, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	129931

**Anti-Nogo Antibody - Additional Information****Gene ID** 57142**Other Names**

KIAA0886; NOGO; Reticulon-4; Foocen; Neurite outgrowth inhibitor; Nogo protein; Neuroendocrine-specific protein; NSP; Neuroendocrine-specific protein C homolog; RTN-x; Reticulon-5

**Target/Specificity**

Recognizes endogenous levels of Nogo protein.

**Dilution**

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200)  
IHC~~1:100~500

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-Nogo Antibody - Protein Information****Name** RTN4 ([HGNC:14085](#))**Function**

Required to induce the formation and stabilization of endoplasmic reticulum (ER) tubules (PubMed: <a href="http://www.uniprot.org/citations/24262037" target="\_blank">24262037</a>, PubMed: <a href="http://www.uniprot.org/citations/25612671" target="\_blank">25612671</a>, PubMed: <a href="http://www.uniprot.org/citations/27619977" target="\_blank">27619977</a>). They regulate membrane morphogenesis in the ER by promoting tubular ER production

(PubMed:<a href="http://www.uniprot.org/citations/24262037" target="\_blank">24262037</a>, PubMed:<a href="http://www.uniprot.org/citations/25612671" target="\_blank">25612671</a>, PubMed:<a href="http://www.uniprot.org/citations/27619977" target="\_blank">27619977</a>, PubMed:<a href="http://www.uniprot.org/citations/27786289" target="\_blank">27786289</a>). They influence nuclear envelope expansion, nuclear pore complex formation and proper localization of inner nuclear membrane proteins (PubMed:<a href="http://www.uniprot.org/citations/26906412" target="\_blank">26906412</a>). However each isoform have specific functions mainly depending on their tissue expression specificities (Probable).

### Cellular Location

[Isoform A]: Endoplasmic reticulum membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein; Cytoplasmic side Synapse {ECO:0000250|UniProtKB:Q99P72}. Note=Anchored to the membrane of the endoplasmic reticulum (ER) through 2 putative transmembrane domains. Localizes throughout the ER tubular network (PubMed:27619977) Co-localizes with TMEM33 at the ER sheets [Isoform C]: Endoplasmic reticulum membrane; Multi-pass membrane protein

### Tissue Location


Isoform A: is specifically expressed in brain and testis and weakly in heart and skeletal muscle. Isoform B: widely expressed except for the liver. Highly expressed in endothelial cells and vascular smooth muscle cells, including blood vessels and mesenteric arteries (PubMed:15034570, PubMed:21183689). Isoform C: is expressed in brain, skeletal muscle and adipocytes. Isoform D is testis-specific.

### Anti-Nogo Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-Nogo Antibody - Images



Western blot analysis of Nogo expression in mouse brain (A), rat brain (B), K562 (C), U87MG (D) whole cell lysates.



Immunohistochemical analysis of Nogo staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **Anti-Nogo Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Nogo. The exact sequence is proprietary.