

### Anti-5-HT1F Antibody

Rabbit polyclonal antibody to 5-HT1F Catalog # AP60467

#### Specification

## Anti-5-HT1F Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Calculated MW WB <u>P30939</u> <u>Q02284</u> Human, Mouse, Rat, Monkey Rabbit Polyclonal 41709

### Anti-5-HT1F Antibody - Additional Information

Gene ID 3355

Other Names HTR1EL; 5-hydroxytryptamine receptor 1F; 5-HT-1F; 5-HT1F; Serotonin receptor 1F

**Target/Specificity** KLH-conjugated synthetic peptide encompassing a sequence within the center region of human 5-HT1F. The exact sequence is proprietary.

Dilution WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

## Anti-5-HT1F Antibody - Protein Information

#### Name HTR1F (HGNC:5292)

Function

G-protein coupled receptor for 5-hydroxytryptamine (serotonin) (PubMed:<a href="http://www.uniprot.org/citations/21422162" target="\_blank">21422162</a>, PubMed:<a href="http://www.uniprot.org/citations/34239069" target="\_blank">34239069</a>, PubMed:<a href="http://www.uniprot.org/citations/8380639" target="\_blank">8380639</a>, PubMed:<a href="http://www.uniprot.org/citations/8380639" target="\_blank">8380639</a>, PubMed:<a href="http://www.uniprot.org/citations/8384716" target="\_blank">8380639</a>, PubMed:<a href="http://www.uniprot.org/citations/8384716" target="\_blank">8384716</a>). Also functions as a receptor for various alkaloids and psychoactive substances (PubMed:<a href="http://www.uniprot.org/citations/21422162" target="\_blank">21422162</a>, PubMed:<a href="http://www.uniprot.org/citations/8380639" target="\_blank">8380639</a>, PubMed:<a



href="http://www.uniprot.org/citations/8384716" target="\_blank">8384716</a>). Receptor for lasmiditan, a drug for the treatment of acute migraine (PubMed:<a

href="http://www.uniprot.org/citations/34239069" target="\_blank">34239069</a>). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors, such as adenylate cyclase (PubMed:<a href="http://www.uniprot.org/citations/34239069"">http://www.uniprot.org/citations/34239069</a>

target="\_blank">34239069</a>). HTR1F is coupled to G(i)/G(o) G alpha proteins and mediates inhibitory neurotransmission by inhibiting adenylate cyclase activity (PubMed:<a href="http://www.uniprot.org/citations/34239069" target="\_blank">34239069</a>, PubMed:<a href="http://www.uniprot.org/citations/35610220" target=" blank">34239069</a>, PubMed:<a href="http://www.uniprot.org/citations/35610220" target=" blank">35610220</a>).

**Cellular Location** 


Cell membrane; Multi-pass membrane protein

## **Anti-5-HT1F Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-5-HT1F Antibody - Images



Western blot analysis of 5-HT1F expression in HEK293T (A), Hela (B), mouse kidney (C), mouse lung (D), mouse liver (E), rat kidney (F), rat lung (G) whole cell lysates.

# Anti-5-HT1F Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human 5-HT1F. The exact sequence is proprietary.