

**Anti-CD61 (pY785) Antibody**  
**Rabbit polyclonal antibody to CD61 (pY785)**  
**Catalog # AP60473**

### Specification

#### Anti-CD61 (pY785) Antibody - Product Information

Application	WB
Primary Accession	<a href="#">P05106</a>
Other Accession	<a href="#">O54890</a>
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Bovine, SARS, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	87058

#### Anti-CD61 (pY785) Antibody - Additional Information

##### Gene ID 3690

##### Other Names

GP3A; Integrin beta-3; Platelet membrane glycoprotein IIIa; GPIIIa; CD61

##### Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human CD61 (pY785). The exact sequence is proprietary.

##### Dilution

WB~~WB (1/500 - 1/1000)

##### Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

##### Storage

Store at -20 °C. Stable for 12 months from date of receipt

#### Anti-CD61 (pY785) Antibody - Protein Information

Name ITGB3 ([HGNC:6156](#))

Synonyms GP3A

##### Function

Integrin alpha-V/beta-3 (ITGAV:ITGB3) is a receptor for cytотactин, fibronectin, laminin, matrix metalloproteinase-2, osteopontin, osteomodulin, prothrombin, thrombospondin, vitronectin and von Willebrand factor. Integrin alpha-IIb/beta-3 (ITGA2B:ITGB3) is a receptor for fibronectin, fibrinogen, plasminogen, prothrombin, thrombospondin and vitronectin. Integrins alpha-IIb/beta-3 and alpha- V/beta-3 recognize the sequence R-G-D in a wide array of ligands. Integrin

alpha-IIb/beta-3 recognizes the sequence H-H-L-G-G-A-K-Q-A- G-D-V in fibrinogen gamma chain (By similarity). Following activation integrin alpha-IIb/beta-3 brings about platelet/platelet interaction through binding of soluble fibrinogen (PubMed:<a href="<http://www.uniprot.org/citations/9111081>">9111081</a>). This step leads to rapid platelet aggregation which physically plugs ruptured endothelial surface. Fibrinogen binding enhances SELP expression in activated platelets (By similarity). ITGAV:ITGB3 binds to fractalkine (CX3CL1) and acts as its coreceptor in CX3CR1-dependent fractalkine signaling (PubMed:<a href="<http://www.uniprot.org/citations/23125415>">23125415</a>, PubMed:<a href="<http://www.uniprot.org/citations/24789099>">24789099</a>). ITGAV:ITGB3 binds to NRG1 (via EGF domain) and this binding is essential for NRG1-ERBB signaling (PubMed:<a href="<http://www.uniprot.org/citations/20682778>">20682778</a>). ITGAV:ITGB3 binds to FGF1 and this binding is essential for FGF1 signaling (PubMed:<a href="<http://www.uniprot.org/citations/18441324>">18441324</a>). ITGAV:ITGB3 binds to FGF2 and this binding is essential for FGF2 signaling (PubMed:<a href="<http://www.uniprot.org/citations/28302677>">28302677</a>). ITGAV:ITGB3 binds to IGF1 and this binding is essential for IGF1 signaling (PubMed:<a href="<http://www.uniprot.org/citations/19578119>">19578119</a>). ITGAV:ITGB3 binds to IGF2 and this binding is essential for IGF2 signaling (PubMed:<a href="<http://www.uniprot.org/citations/28873464>">28873464</a>). ITGAV:ITGB3 binds to IL1B and this binding is essential for IL1B signaling (PubMed:<a href="<http://www.uniprot.org/citations/29030430>">29030430</a>). ITGAV:ITGB3 binds to PLA2G2A via a site (site 2) which is distinct from the classical ligand-binding site (site 1) and this induces integrin conformational changes and enhanced ligand binding to site 1 (PubMed:<a href="<http://www.uniprot.org/citations/18635536>">18635536</a>, PubMed:<a href="<http://www.uniprot.org/citations/25398877>">25398877</a>). ITGAV:ITGB3 acts as a receptor for fibrillin-1 (FBN1) and mediates R-G-D-dependent cell adhesion to FBN1 (PubMed:<a href="<http://www.uniprot.org/citations/12807887>">12807887</a>). In brain, plays a role in synaptic transmission and plasticity. Involved in the regulation of the serotonin neurotransmission, is required to localize to specific compartments within the synapse the serotonin receptor SLC6A4 and for an appropriate reuptake of serotonin. Controls excitatory synaptic strength by regulating GRIA2-containing AMPAR endocytosis, which affects AMPAR abundance and composition (By similarity). ITGAV:ITGB3 act as a receptor for CD40LG (PubMed:<a href="<http://www.uniprot.org/citations/31331973>">31331973</a>). ITGAV:ITGB3 acts as a receptor for IBSP and promotes cell adhesion and migration to IBSP (PubMed:<a href="<http://www.uniprot.org/citations/10640428>">10640428</a>).

### Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell projection, lamellipodium membrane. Cell junction, focal adhesion. Postsynaptic cell membrane {ECO:0000250|UniProtKB:O54890}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:O54890}. Synapse {ECO:0000250|UniProtKB:O54890}

### Tissue Location

Isoform beta-3A and isoform beta-3C are widely expressed. Isoform beta-3A is specifically expressed in osteoblast cells; isoform beta-3C is specifically expressed in prostate and testis


### Anti-CD61 (pY785) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-CD61 (pY785) Antibody - Images



Western blot analysis of CD61 (pY785) expression in U87MG (A), PC3 (B) whole cell lysates.



Direct ELISA antibody dose-response curve using Anti-CD61 (pY785) Antibody. Antigen (phosphopeptide and non-phosphopeptide) concentration is 5 ug/ml. Goat Anti-Rabbit IgG (H&L) - HRP was used as the secondary antibody, and signal was developed by TMB substrate.

### Anti-CD61 (pY785) Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human CD61 (pY785). The exact sequence is proprietary.