

Anti-NSG1 Antibody

Rabbit polyclonal antibody to NSG1 Catalog # AP60601

Specification

Anti-NSG1 Antibody - Product Information

Application WB, IHC
Primary Accession P42857
Other Accession O62092

Reactivity Human, Mouse, Rat, Monkey Rabbit

Clonality Polyclonal Calculated MW 20913

Anti-NSG1 Antibody - Additional Information

Gene ID 27065

Other Names

D4S234; Neuron-specific protein family member 1; Brain neuron cytoplasmic protein 1

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NSG1. The exact sequence is proprietary.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200) IHC~~1:100~500

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

Anti-NSG1 Antibody - Protein Information

Name NSG1 (HGNC:18790)

Function

Plays a role in the recycling mechanism in neurons of multiple receptors, including AMPAR, APP and L1CAM and acts at the level of early endosomes to promote sorting of receptors toward a recycling pathway. Regulates sorting and recycling of GRIA2 through interaction with GRIP1 and then contributes to the regulation of synaptic transmission and plasticity by affecting the recycling and targeting of AMPA receptors to the synapse (By similarity). Is required for faithful sorting of L1CAM to axons by facilitating trafficking from somatodendritic early endosome or the recycling endosome (By similarity). In an other hand, induces apoptosis via the activation of CASP3 in



response to DNA damage (PubMed:20599942, PubMed:20878061).

Cellular Location


Membrane {ECO:0000250|UniProtKB:P02683}; Single- pass type II membrane protein {ECO:0000250|UniProtKB:P02683}. Golgi apparatus, trans-Golgi network membrane {ECO:0000250|UniProtKB:P02683} Endosome membrane {ECO:0000250|UniProtKB:P02683}. Cell projection, dendrite {ECO:0000250|UniProtKB:P02683}. Early endosome membrane {ECO:0000250|UniProtKB:P02683}. Late endosome membrane {ECO:0000250|UniProtKB:P02683}. Lysosome lumen {ECO:0000250|UniProtKB:P02683}. Recycling endosome membrane {ECO:0000250|UniProtKB:P02683}. Golgi apparatus, Golgi stack membrane {ECO:0000250|UniProtKB:P02683}. Endosome, multivesicular body membrane {ECO:0000250|UniProtKB:P02683}. Endosome, multivesicular body membrane {ECO:0000250|UniProtKB:P02683}. Endoplasmic reticulum membrane. Note=Endocytosed from the cell surface, thus enters into early endosomes, trafficks to late endosomes and degradates in lysosomes (By similarity). Endoplasmic reticulum targeting is essential for apoptosis (PubMed:20599942). Found in both stationary and motile endosomes. A previous study supports a type I membrane protein topology (By similarity) {ECO:0000250|UniProtKB:P02683, ECO:0000250|UniProtKB:Q62092, ECO:0000269|PubMed:20599942}

Anti-NSG1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-NSG1 Antibody - Images



Western blot analysis of NSG1 expression in mouse muscle (A), rat muscle (B) whole cell lysates.





Immunohistochemical analysis of NSG1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-NSG1 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NSG1. The exact sequence is proprietary.