

# **Anti-PEA15 Antibody**

Rabbit polyclonal antibody to PEA15 Catalog # AP60606

## **Specification**

# **Anti-PEA15 Antibody - Product Information**

Application
Primary Accession
Other Accession
Reactivity
Host

Clonality Calculated MW WB, IP, IF/IC, IHC

<u>Q15121</u> <u>Q62048</u>

Human, Mouse, Rat

Rabbit Polyclonal 15040

# **Anti-PEA15 Antibody - Additional Information**

### **Gene ID 8682**

### **Other Names**

Astrocytic phosphoprotein PEA-15; 15 kDa phosphoprotein enriched in astrocytes; Phosphoprotein enriched in diabetes; PED

# **Target/Specificity**

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human PEA15. The exact sequence is proprietary.

#### **Dilution**

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500), IP (1/10 - 1/100) IP~~N/A IF/IC~~N/A IHC~~1:100~500

#### Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

# Storage


Store at -20 °C.Stable for 12 months from date of receipt


# **Anti-PEA15 Antibody - Protein Information**

### Name PEA15

#### **Function**

Blocks Ras-mediated inhibition of integrin activation and modulates the ERK MAP kinase cascade. Inhibits RPS6KA3 activities by retaining it in the cytoplasm (By similarity). Inhibits both TNFRSF6-and TNFRSF1A-mediated CASP8 activity and apoptosis. Regulates glucose transport by controlling both the content of SLC2A1 glucose transporters on the plasma membrane and the







insulin-dependent trafficking of SLC2A4 from the cell interior to the surface.

### **Cellular Location**

Cytoplasm. Note=Associated with microtubules.

### **Tissue Location**


Ubiquitously expressed. Most abundant in tissues such as heart, brain, muscle and adipose tissue which utilize glucose as an energy source. Lower expression in glucose-producing tissues Higher levels of expression are found in tissues from individuals with type 2 diabetes than in controls.

# **Anti-PEA15 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## Anti-PEA15 Antibody - Images




Western blot analysis of PEA15 expression in mouse brain (A), C6 (B), HEK293T (C) whole cell lysates.





Immunohistochemical analysis of PEA15 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of PEA15 staining in MDA-MB-231 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

### Anti-PEA15 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human PEA15. The exact sequence is proprietary.