

Anti-SAP49 Antibody
Rabbit polyclonal antibody to SAP49
Catalog # AP60631**Specification**

Anti-SAP49 Antibody - Product Information

Application	WB, IHC
Primary Accession	Q15427
Other Accession	Q8QZY9
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44386

Anti-SAP49 Antibody - Additional Information**Gene ID** 10262**Other Names**

SAP49; Splicing factor 3B subunit 4; Pre-mRNA-splicing factor SF3b 49 kDa subunit; SF3b50; Spliceosome-associated protein 49; SAP 49

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human SAP49. The exact sequence is proprietary.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200)

IHC~~1:100~500

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

Anti-SAP49 Antibody - Protein Information**Name** SF3B4**Synonyms** SAP49**Function**

Component of the 17S U2 SnRNP complex of the spliceosome, a large ribonucleoprotein complex that removes introns from transcribed pre-mRNAs (PubMed:10882114, PubMed:12234937, PubMed:12234937, PubMed:12234937)

[27720643](http://www.uniprot.org/citations/27720643), PubMed: [32494006](http://www.uniprot.org/citations/32494006)). The 17S U2 SnRNP complex (1) directly participates in early spliceosome assembly and (2) mediates recognition of the intron branch site during pre-mRNA splicing by promoting the selection of the pre-mRNA branch-site adenosine, the nucleophile for the first step of splicing (PubMed: [12234937](http://www.uniprot.org/citations/12234937), PubMed: [32494006](http://www.uniprot.org/citations/32494006)). Within the 17S U2 SnRNP complex, SF3B4 is part of the SF3B subcomplex, which is required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpoint sequence in pre-mRNA (PubMed: [12234937](http://www.uniprot.org/citations/12234937), PubMed: [27720643](http://www.uniprot.org/citations/27720643)). Sequence independent binding of SF3A and SF3B subcomplexes upstream of the branch site is essential, it may anchor U2 snRNP to the pre-mRNA (PubMed: [12234937](http://www.uniprot.org/citations/12234937)). May also be involved in the assembly of the 'E' complex (PubMed: [10882114](http://www.uniprot.org/citations/10882114)). Also acts as a component of the minor spliceosome, which is involved in the splicing of U12-type introns in pre-mRNAs (PubMed: [15146077](http://www.uniprot.org/citations/15146077), PubMed: [33509932](http://www.uniprot.org/citations/33509932)).

Cellular Location


Nucleus

Anti-SAP49 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-SAP49 Antibody - Images



Western blot analysis of SAP49 expression in LO2 (A), A549 (B), HEK293T (C), CT26 (D), PC12 (E)

whole cell lysates.



Immunohistochemical analysis of SAP49 staining in human lung formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-SAP49 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human SAP49. The exact sequence is proprietary.