

**Anti-GIP Receptor Antibody**  
**Rabbit polyclonal antibody to GIP Receptor**  
**Catalog # AP60804****Specification**

---

**Anti-GIP Receptor Antibody - Product Information**

Application	WB, IF/IC, IHC
Primary Accession	<a href="#">P48546</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53157

**Anti-GIP Receptor Antibody - Additional Information****Gene ID** 2696**Other Names**

Gastric inhibitory polypeptide receptor; GIP-R; Glucose-dependent insulinotropic polypeptide receptor

**Target/Specificity**

Recognizes endogenous levels of GIP Receptor protein.

**Dilution**

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500)

IF/IC~~N/A

IHC~~1:100~500

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-GIP Receptor Antibody - Protein Information****Name** GIPR**Function**

This is a receptor for GIP. The activity of this receptor is mediated by G proteins which activate adenylyl cyclase.

**Cellular Location**


Cell membrane; Multi-pass membrane protein

## Anti-GIP Receptor Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## Anti-GIP Receptor Antibody - Images



Western blot analysis of GIP Receptor expression in HepG2 (A) whole cell lysates.



Immunohistochemical analysis of GIP Receptor staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of GIP Receptor staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

#### **Anti-GIP Receptor Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human GIP Receptor. The exact sequence is proprietary.