

Anti-AXL Antibody

Rabbit polyclonal antibody to AXL Catalog # AP60816

Specification

Anti-AXL Antibody - Product Information

Application
Primary Accession
Other Accession
Reactivity
Host
Clonality
Calculated MW

WB, IHC
P30530
O00993
Human, Mouse, Rat
Rabbit
Polyclonal
98337

Anti-AXL Antibody - Additional Information

Gene ID 558

Other Names

UFO; Tyrosine-protein kinase receptor UFO; AXL oncogene

Target/Specificity

Recognizes endogenous levels of AXL protein.

Dilution

WB~~WB (1/500 - 1/2000), IH (1/50 - 1/200) IHC~~1:100~500

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-AXL Antibody - Protein Information

Name AXL

Synonyms UFO

Function

Receptor tyrosine kinase that transduces signals from the extracellular matrix into the cytoplasm by binding growth factor GAS6 and which is thus regulating many physiological processes including cell survival, cell proliferation, migration and differentiation. Ligand binding at the cell surface induces dimerization and autophosphorylation of AXL. Following activation by ligand, AXL binds and induces tyrosine phosphorylation of PI3-kinase subunits PIK3R1, PIK3R2 and PIK3R3; but also GRB2, PLCG1, LCK and PTPN11. Other downstream substrate candidates for AXL are CBL,



NCK2, SOCS1 and TNS2. Recruitment of GRB2 and phosphatidylinositol 3 kinase regulatory subunits by AXL leads to the downstream activation of the AKT kinase. GAS6/AXL signaling plays a role in various processes such as endothelial cell survival during acidification by preventing apoptosis, optimal cytokine signaling during human natural killer cell development, hepatic regeneration, gonadotropin-releasing hormone neuron survival and migration, platelet activation, or regulation of thrombotic responses. Also plays an important role in inhibition of Toll-like

Cellular Location

Cell membrane; Single-pass type I membrane protein

receptors (TLRs)-mediated innate immune response.

Tissue Location


Highly expressed in metastatic colon tumors. Expressed in primary colon tumors. Weakly expressed in normal colon tissue.

Anti-AXL Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-AXL Antibody - Images



Western blot analysis of AXL expression in PC3 (A), HEK293T (B), U87MG (C), SKOVCAR3 (D), C6 (E), CT26 (F) whole cell lysates.





Immunohistochemical analysis of AXL staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-AXL Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human AXL. The exact sequence is proprietary.