

**Anti-ALPK1 Antibody**  
**Rabbit polyclonal antibody to ALPK1**  
**Catalog # AP60840****Specification**

---

**Anti-ALPK1 Antibody - Product Information**

Application	WB, IF/IC, IHC
Primary Accession	<a href="#">Q96QP1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	138861

**Anti-ALPK1 Antibody - Additional Information****Gene ID** 80216**Other Names**

KIAA1527; LAK; Alpha-protein kinase 1; Chromosome 4 kinase; Lymphocyte alpha-protein kinase

**Target/Specificity**

Recognizes endogenous levels of ALPK1 protein.

**Dilution**

WB~~WB (1/500 - 1/2000), IH (1/50 - 1/200), IF/IC (1/50 - 1/100)

IF/IC~~N/A

IHC~~1:100~500

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-ALPK1 Antibody - Protein Information****Name** ALPK1 {ECO:0000303|PubMed:30111836, ECO:0000312|HGNC:HGNC:20917}**Function**

Serine/threonine-protein kinase that detects bacterial pathogen-associated molecular pattern metabolites (PAMPs) and initiates an innate immune response, a critical step for pathogen elimination and engagement of adaptive immunity (PubMed:<a href="http://www.uniprot.org/citations/28222186" target="\_blank">28222186</a>, PubMed:<a href="http://www.uniprot.org/citations/28877472" target="\_blank">28877472</a>, PubMed:<a href="http://www.uniprot.org/citations/30111836" target="\_blank">30111836</a>). Specifically recognizes and binds ADP-D-glycero-beta- D-manno-heptose (ADP-Heptose), a potent PAMP present in all Gram- negative and some Gram-positive bacteria (PubMed:<a

[30111836](http://www.uniprot.org/citations/30111836)). ADP-Heptose-binding stimulates its kinase activity to phosphorylate and activate TIFA, triggering pro-inflammatory NF-kappa-B signaling (PubMed:[30111836](http://www.uniprot.org/citations/30111836)). May be involved in monosodium urate monohydrate (MSU)-induced inflammation by mediating phosphorylation of unconventional myosin MYO9A (PubMed:[27169898](http://www.uniprot.org/citations/27169898)). May also play a role in apical protein transport by mediating phosphorylation of unconventional myosin MYO1A (PubMed:[15883161](http://www.uniprot.org/citations/15883161)). May play a role in ciliogenesis (PubMed:[30967659](http://www.uniprot.org/citations/30967659)).

#### Cellular Location

Cytoplasm, cytosol. Cytoplasm, cytoskeleton, spindle pole Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cell projection, cilium. Note=Localized at the base of primary cilia.

#### Tissue Location


Highly expressed in liver. Expressed in the optic nerve and retinal pigmented epithelium. Lower expression is observed in the macula and extramacular retina (PubMed:30967659)

### Anti-ALPK1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### Anti-ALPK1 Antibody - Images



Western blot analysis of ALPK1 expression in HEK293T (A) whole cell lysates.



Immunohistochemical analysis of ALPK1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of ALPK1 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

#### **Anti-ALPK1 Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human ALPK1. The exact sequence is proprietary.