

## **Anti-UCP2 Antibody**

Rabbit polyclonal antibody to UCP2 Catalog # AP60867

## **Specification**

## **Anti-UCP2 Antibody - Product Information**

Application WB, IHC
Primary Accession P55851
Other Accession P70406

Reactivity Human, Mouse, Rat, Bovine, SARS, Dog Rabbit

Clonality Polyclonal Calculated MW 33229

## **Anti-UCP2 Antibody - Additional Information**

#### **Gene ID 7351**

#### **Other Names**

SLC25A8; Mitochondrial uncoupling protein 2; UCP 2; Solute carrier family 25 member 8; UCPH

## Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human UCP2. The exact sequence is proprietary.

#### Dilution

WB~~WB (1/500 - 1/1000), IH (1/50 - 1/100) IHC~~1:100~500

#### **Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

## **Storage**

Store at -20 °C.Stable for 12 months from date of receipt

## **Anti-UCP2 Antibody - Protein Information**

## Name UCP2

**Synonyms** SLC25A8 {ECO:0000303|PubMed:33798544}

#### **Function**

Antiporter that exports dicarboxylate intermediates of the Krebs cycle in exchange for phosphate plus a proton across the inner membrane of mitochondria, a process driven by mitochondrial motive force with an overall impact on glycolysis, glutaminolysis and glutathione-dependent redox balance. Continuous export of oxaloacetate and related four-carbon dicarboxylates from mitochondrial matrix into the cytosol negatively regulates the oxidation of acetyl-CoA substrates



via the Krebs cycle, lowering the ATP/ADP ratio and reactive oxygen species (ROS) production (PubMed:<a href="http://www.uniprot.org/citations/24395786" target="\_blank">24395786</a>). May mediate inducible proton entry into the mitochondrial matrix affecting ATP turnover as a protection mechanism against oxidative stress. The proton currents are most likely associated with fatty acid flipping across the inner membrane of mitochondria in a metabolic process regulated by free fatty acids and purine nucleotides (By similarity) (PubMed:<a

href="http://www.uniprot.org/citations/11171965" target=" blank">11171965</a>, PubMed:<a href="http://www.uniprot.org/citations/11278935" target="blank">11278935</a>, PubMed:<a href="http://www.uniprot.org/citations/22524567" target="blank">22524567</a>, PubMed:<a href="http://www.uniprot.org/citations/26182433" target="\_blank">26182433</a>, PubMed:<a href="http://www.uniprot.org/citations/33373220" target="blank">33373220</a>). Regulates the use of glucose as a source of energy. Required for glucose-induced DRP1-dependent mitochondrial fission and neuron activation in the ventromedial nucleus of the hypothalamus (VMH). This mitochondrial adaptation mechanism modulates the VMH pool of glucose- excited neurons with an impact on systemic glucose homeostasis (By similarity). Regulates ROS levels and metabolic reprogramming of macrophages during the resolution phase of inflammation. Attenuates ROS production in response to IL33 to preserve the integrity of the Krebs cycle required for persistent production of itaconate and subsequent GATA3-dependent differentiation of inflammation-resolving alternatively activated macrophages (By similarity). Can unidirectionally transport anions including L-malate, L-aspartate, phosphate and chloride ions (PubMed: <a href="http://www.uniprot.org/citations/22524567" target=" blank">22524567</a>, PubMed:<a href="http://www.uniprot.org/citations/24395786" target="\_blank">24395786</a>, PubMed:<a href="http://www.uniprot.org/citations/26182433" target="blank">26182433</a>). Does not mediate adaptive thermogenesis (By similarity).

#### **Cellular Location**

Mitochondrion inner membrane {ECO:0000250|UniProtKB:P70406}; Multi-pass membrane protein

#### **Tissue Location**

Widely expressed in adult human tissues, including tissues rich in macrophages. Most expressed in white adipose tissue and skeletal muscle.


## **Anti-UCP2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# **Anti-UCP2 Antibody - Images**





Western blot analysis of UCP2 expression in HEK293T (A), Raw264.7 (B), PC12 (C) whole cell lysates.



Immunohistochemical analysis of UCP2 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## Anti-UCP2 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human UCP2. The exact sequence is proprietary.