

Anti-USP28 Antibody

Rabbit polyclonal antibody to USP28 Catalog # AP60942

Specification

Anti-USP28 Antibody - Product Information

Application WB
Primary Accession Q96RU2
Other Accession Q51043
Reactivity Human, Mouse

Host Rabbit
Clonality Polyclonal
Calculated MW 122491

Anti-USP28 Antibody - Additional Information

Gene ID 57646

Other Names

KIAA1515; Ubiquitin carboxyl-terminal hydrolase 28; Deubiquitinating enzyme 28; Ubiquitin thioesterase 28; Ubiquitin-specific-processing protease 28

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human USP28. The exact sequence is proprietary.

Dilution

WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-USP28 Antibody - Protein Information

Name USP28

Synonyms KIAA1515

Function

Deubiquitinase involved in DNA damage response checkpoint and MYC proto-oncogene stability. Involved in DNA damage induced apoptosis by specifically deubiquitinating proteins of the DNA damage pathway such as CLSPN. Also involved in G2 DNA damage checkpoint, by deubiquitinating CLSPN, and preventing its degradation by the anaphase promoting complex/cyclosome (APC/C). In contrast, it does not deubiquitinate PLK1. Specifically deubiquitinates MYC in the nucleoplasm,



leading to prevent MYC degradation by the proteasome: acts by specifically interacting with isoform 1 of FBXW7 (FBW7alpha) in the nucleoplasm and counteracting ubiquitination of MYC by the SCF(FBW7) complex. In contrast, it does not interact with isoform 4 of FBXW7 (FBW7gamma) in the nucleolus, allowing MYC degradation and explaining the selective MYC degradation in the nucleolus. Deubiquitinates ZNF304, hence preventing ZNF304 degradation by the proteasome and leading to the activated KRAS-mediated promoter hypermethylation and transcriptional silencing of tumor suppressor genes (TSGs) in a subset of colorectal cancers (CRC) cells (PubMed:24623306).


Cellular Location Nucleus, nucleoplasm

Anti-USP28 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-USP28 Antibody - Images



Western blot analysis of USP28 expression in HEK293T (A), A549 (B), mouse heart (C) whole cell lysates.

Anti-USP28 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human USP28. The exact sequence is proprietary.