

Anti-IFNAR2 Antibody
Rabbit polyclonal antibody to IFNAR2
Catalog # AP61130

Specification

Anti-IFNAR2 Antibody - Product Information

Application	WB, IH
Primary Accession	P48551
Other Accession	O35664
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57759

Anti-IFNAR2 Antibody - Additional Information

Gene ID 3455

Other Names

IFNABR; IFNARB; Interferon alpha/beta receptor 2; IFN-R-2; IFN-alpha binding protein; IFN-alpha/beta receptor 2; Interferon alpha binding protein; Type I interferon receptor 2

Target/Specificity

Recognizes endogenous levels of IFNAR2 protein.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/50 - 1/200)
IH~~WB (1/500 - 1/1000), IH (1/50 - 1/200)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-IFNAR2 Antibody - Protein Information

Name IFNAR2

Synonyms IFNABR, IFNARB

Function

Together with IFNAR1, forms the heterodimeric receptor for type I interferons (including interferons alpha, beta, epsilon, omega and kappa) (PubMed:8181059, PubMed:7665574, PubMed:7759950, PubMed:11111111)

href="http://www.uniprot.org/citations/8798579" target="_blank">>8798579, PubMed:>8969169, PubMed:>10049744, PubMed:>10556041, PubMed:>21854986, PubMed:>26424569, PubMed:>28165510, PubMed:>32972995). Type I interferon binding activates the JAK-STAT signaling cascade, resulting in transcriptional activation or repression of interferon- regulated genes that encode the effectors of the interferon response (PubMed:>8181059, PubMed:>7665574, PubMed:>7759950, PubMed:>8798579, PubMed:>8969169, PubMed:>10049744, PubMed:>17517919, PubMed:>21854986, PubMed:>26424569, PubMed:>28165510, PubMed:>32972995). Mechanistically, type I interferon-binding brings the IFNAR1 and IFNAR2 subunits into close proximity with one another, driving their associated Janus kinases (JAKs) (TYK2 bound to IFNAR1 and JAK1 bound to IFNAR2) to cross-phosphorylate one another (PubMed:>10556041, PubMed:>11682488, PubMed:>12105218, PubMed:>21854986, PubMed:>32972995). The activated kinases phosphorylate specific tyrosine residues on the intracellular domains of IFNAR1 and IFNAR2, forming docking sites for the STAT transcription factors (STAT1, STAT2 and STAT) (PubMed:>11682488, PubMed:>12105218, PubMed:>21854986, PubMed:>32972995). STAT proteins are then phosphorylated by the JAKs, promoting their translocation into the nucleus to regulate expression of interferon-regulated genes (PubMed:>9121453, PubMed:>12105218, PubMed:>28165510).

Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein [Isoform 3]: Secreted

Tissue Location


Isoform 3 is detected in the urine (at protein level) (PubMed:8181059, PubMed:7759950). Expressed in blood cells Expressed in lymphoblastoid and fibrosarcoma cell lines

Anti-IFNAR2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IFNAR2 Antibody - Images

Western blot analysis of IFNAR2 expression in PC12 (A), NIH3T3 (B), MCF7 (C), PC3 (D) whole cell lysates.



Immunohistochemical analysis of IFNAR2 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-IFNAR2 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human IFNAR2. The exact sequence is proprietary.