

### Anti-GPR106 Antibody

Rabbit polyclonal antibody to GPR106 Catalog # AP61358

#### Specification

## **Anti-GPR106 Antibody - Product Information**

Application Primary Accession Other Accession Reactivity Host Clonality Calculated MW WB, IF/IC <u>O8WXD0</u> <u>O91ZZ5</u> Human, Mouse, Dog Rabbit Polyclonal 86453

### **Anti-GPR106 Antibody - Additional Information**

Gene ID 122042

**Other Names** GPR106; GREAT; LGR8; Relaxin receptor 2; G-protein coupled receptor 106; G-protein coupled receptor affecting testicular descent; Leucine-rich repeat-containing G-protein coupled receptor 8; Relaxin family peptide receptor 2

Target/Specificity Recognizes endogenous levels of GPR106 protein.

Dilution WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500) IF/IC~~N/A

Format Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

### **Anti-GPR106 Antibody - Protein Information**

Name RXFP2

Synonyms GPR106, GREAT, LGR8

Function

Receptor for relaxin. The activity of this receptor is mediated by G proteins leading to stimulation of adenylate cyclase and an increase of cAMP. May also be a receptor for Leydig insulin-like peptide (INSL3).



**Cellular Location** 

Cell membrane; Multi-pass membrane protein.

**Tissue Location** 

Expressed mainly in the brain, kidney, muscle, testis, thyroid, uterus, peripheral blood cells and bone marrow

# **Anti-GPR106 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

## Anti-GPR106 Antibody - Images



Western blot analysis of GPR106 expression in Hela (A), Jurkat (B), mouse brain (C) whole cell lysates.



Immunofluorescent analysis of GPR106 staining in NIH3T3 cells. Formalin-fixed cells were



permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

### Anti-GPR106 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human GPR106. The exact sequence is proprietary.