

**Anti-FGF1 Antibody**  
**Rabbit polyclonal antibody to FGF1**  
**Catalog # AP61406****Specification**

---

**Anti-FGF1 Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">P05230</a>
Other Accession	<a href="#">P61148</a>
Reactivity	Human, Mouse, Rat, Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17460

**Anti-FGF1 Antibody - Additional Information****Gene ID** 2246**Other Names**

FGFA; Fibroblast growth factor 1; FGF-1; Acidic fibroblast growth factor; aFGF; Endothelial cell growth factor; ECGF; Heparin-binding growth factor 1; HBGF-1

**Target/Specificity**

Recognizes endogenous levels of FGF1 protein.

**Dilution**WB~~WB (1/500 - 1/1000), IH (1/50 - 1/200)  
IHC~~1:100~500**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C.Stable for 12 months from date of receipt

**Anti-FGF1 Antibody - Protein Information****Name** FGF1**Synonyms** FGFA**Function**

Plays an important role in the regulation of cell survival, cell division, angiogenesis, cell differentiation and cell migration. Functions as a potent mitogen in vitro. Acts as a ligand for FGFR1 and integrins. Binds to FGFR1 in the presence of heparin leading to FGFR1 dimerization and activation via sequential autophosphorylation on tyrosine residues which act as docking sites for interacting proteins, leading to the activation of several signaling cascades. Binds to integrin

ITGAV:ITGB3. Its binding to integrin, subsequent ternary complex formation with integrin and FGFR1, and the recruitment of PTPN11 to the complex are essential for FGF1 signaling. Induces the phosphorylation and activation of FGFR1, FRS2, MAPK3/ERK1, MAPK1/ERK2 and AKT1 (PubMed:<a href="http://www.uniprot.org/citations/18441324" target="\_blank">18441324</a>, PubMed:<a href="http://www.uniprot.org/citations/20422052" target="\_blank">20422052</a>). Can induce angiogenesis (PubMed:<a href="http://www.uniprot.org/citations/23469107" target="\_blank">23469107</a>).

#### Cellular Location

Secreted. Cytoplasm. Cytoplasm, cell cortex. Cytoplasm, cytosol. Nucleus. Note=Lacks a cleavable signal sequence Within the cytoplasm, it is transported to the cell membrane and then secreted by a non-classical pathway that requires Cu(2+) ions and S100A13. Secreted in a complex with SYT1 (By similarity). Binding of exogenous FGF1 to FGFR facilitates endocytosis followed by translocation of FGF1 across endosomal membrane into the cytosol Nuclear import from the cytosol requires the classical nuclear import machinery, involving proteins KPNA1 and KPNB1, as well as LRRC59

#### Tissue Location


Predominantly expressed in kidney and brain. Detected at much lower levels in heart and skeletal muscle

### Anti-FGF1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-FGF1 Antibody - Images



Western blot analysis of FGF1 expression in H1688 (A), mouse kidney (B), mouse brain (C), rat kidney (D), rat brain (E) whole cell lysates.



Immunohistochemical analysis of FGF1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

**Anti-FGF1 Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human FGF1. The exact sequence is proprietary.