

Anti-Histone H4 (MonoMethyl-K5) Antibody
Rabbit polyclonal antibody to Histone H4 (MonoMethyl-K5)
Catalog # AP61436**Specification**

Anti-Histone H4 (MonoMethyl-K5) Antibody - Product Information

Application	WB, IH, E
Primary Accession	P62805
Other Accession	P62806
Reactivity	Human, Mouse, Rat, Pig, Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	11367

Anti-Histone H4 (MonoMethyl-K5) Antibody - Additional Information

Gene ID 121504;554313;8294;8359;8360;8361;8362;8363;8364;8365;8366;8367;8368;8370

Other Names

H4/A; H4FA; H4/I; H4FI; H4/G; H4FG; H4/B; H4FB; H4/J; H4FJ; H4/C; H4FC; H4/H; H4FH; H4/M; H4FM; H4/E; H4FE; H4/D; H4FD; H4/K; H4FK; H4/N; H4F2; H4FN; HIST2H4; H4/O; H4FO; Histone H4

Target/Specificity

Recognizes endogenous levels of Histone H4 with a site at MonoMethyl-K5 protein.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/50 - 1/200)

IH~~WB (1/500 - 1/1000), IH (1/50 - 1/200)

E~~WB (1/500 - 1/1000), IH (1/50 - 1/200)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-Histone H4 (MonoMethyl-K5) Antibody - Protein Information

Name H4C1

Synonyms H4/A, H4FA, HIST1H4A

Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of

histones, also called histone code, and nucleosome remodeling.

Cellular Location


Nucleus. Chromosome.

Anti-Histone H4 (MonoMethyl-K5) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Histone H4 (MonoMethyl-K5) Antibody - Images




Western blot analysis of Histone H4 (MonoMethyl-K5) expression in HeLa (A) whole cell lysates.



Immunohistochemical analysis of Histone H4 (MonoMethyl-K5) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system.

DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Direct ELISA antibody dose-response curve using Anti-Histone H4 (MonoMethyl-K5) Antibody. Antigen (methyl-peptide and non-methyl-peptide) concentration is 5 ug/ml. Goat Anti-Rabbit IgG (H&L) - HRP was used as the secondary antibody, and signal was developed by TMB substrate.

Anti-Histone H4 (MonoMethyl-K5) Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Histone H4 with a site at MonoMethyl-K5. The exact sequence is proprietary.