

Anti-Defensin alpha 3 Antibody

Rabbit polyclonal antibody to Defensin alpha 3 Catalog # AP61481

Specification

Anti-Defensin alpha 3 Antibody - Product Information

Application Primary Accession	WB, IHC P59666
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	10245

Anti-Defensin alpha 3 Antibody - Additional Information

Gene ID 1668

Other Names DEF3; Neutrophil defensin 3; Defensin alpha 3; HNP-3; HP-3; HP3

Target/Specificity Recognizes endogenous levels of Defensin alpha 3 protein.

Dilution WB~~WB (1/500 - 1/1000), IH (1/50 - 1/200) IHC~~1:100~500

Format Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

Anti-Defensin alpha 3 Antibody - Protein Information

Name DEFA3

Synonyms DEF3

Function

Effector molecule of the innate immune system that acts via antibiotic-like properties against a broad array of infectious agents including bacteria, fungi, and viruses (PubMed:15616305, PubMed:15772169, PubMed:15772169, PubMed:17142766, PubMed:17142766). Possesses the ability to neutralize bacterial toxins such as B.anthracis lethal factor, Clostridium difficile cytotoxin B as well as leukocidin produced by Staphylococcus aureus (PubMed:<a



href="http://www.uniprot.org/citations/15772169" target="_blank">15772169, PubMed:18435932, PubMed:25963798). Also blocks
herpes simplex virus infection by interacting with envelope glycoprotein B and thus preventing its
binding to heparan sulfate, the receptor for attachment (PubMed:17142766).


Cellular Location Secreted.

Anti-Defensin alpha 3 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Defensin alpha 3 Antibody - Images



Western blot analysis of Defensin alpha 3 expression in H1688 (A) whole cell lysates.





Immunohistochemical analysis of Defensin alpha 3 staining in human skeletal muscle formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-Defensin alpha 3 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Defensin alpha 3. The exact sequence is proprietary.