

Anti-Pyruvate Kinase Antibody
Rabbit polyclonal antibody to Pyruvate Kinase
Catalog # AP61538**Specification**

Anti-Pyruvate Kinase Antibody - Product Information

Application	WB
Primary Accession	P14618
Other Accession	P52480
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57937

Anti-Pyruvate Kinase Antibody - Additional Information**Gene ID** 5315**Other Names**

OIP3; PK2; PK3; PKM2; Pyruvate kinase PKM; Cytosolic thyroid hormone-binding protein; CTHBP; Opa-interacting protein 3; OIP-3; Pyruvate kinase 2/3; Pyruvate kinase muscle isozyme; Thyroid hormone-binding protein 1; THBP1; Tumor M2-PK; p58

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Pyruvate Kinase. The exact sequence is proprietary.

Dilution

WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-Pyruvate Kinase Antibody - Protein Information**Name** PKM**Synonyms** OIP3 {ECO:0000303|PubMed:9466265}, PK2,**Function**

Catalyzes the final rate-limiting step of glycolysis by mediating the transfer of a phosphoryl group from phosphoenolpyruvate (PEP) to ADP, generating ATP (PubMed:15996096, PubMed:1854723, PubMed:20847263). The ratio between the highly active tetrameric form and nearly inactive dimeric form determines whether glucose carbons are channeled to biosynthetic processes or used for glycolytic ATP production (PubMed:15996096, PubMed:1854723, PubMed:20847263). The transition between the 2 forms contributes to the control of glycolysis and is important for tumor cell proliferation and survival (PubMed:15996096, PubMed:1854723, PubMed:20847263).

Cellular Location

[Isoform M2]: Cytoplasm. Nucleus Note=Translocates to the nucleus in response to various signals, such as EGF receptor activation or apoptotic stimuli (PubMed:17308100, PubMed:22056988, PubMed:24120661). Nuclear translocation is promoted by acetylation by EP300 (PubMed:24120661). Deacetylation by SIRT6 promotes its nuclear export in a process dependent of XPO4, thereby suppressing its ability to activate transcription and promote tumorigenesis (PubMed:26787900).

Tissue Location


[Isoform M2]: Specifically expressed in proliferating cells, such as embryonic stem cells, embryonic carcinoma cells, as well as cancer cells.

Anti-Pyruvate Kinase Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Pyruvate Kinase Antibody - Images



Western blot analysis of Pyruvate Kinase expression in MCF7 (A), HepG2 (B) whole cell lysates.

Anti-Pyruvate Kinase Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Pyruvate Kinase. The exact sequence is proprietary.