

Anti-Myoglobin Antibody

Mouse monoclonal antibody to Myoglobin Catalog # AP61624

Specification

Anti-Myoglobin Antibody - Product Information

Application IHC
Primary Accession P02144
Reactivity Human
Host Mouse
Clonality Monoclonal
Calculated MW 17184

Anti-Myoglobin Antibody - Additional Information

Gene ID 4151

Other Names Myoglobin

Target/Specificity

Recognizes endogenous levels of Myoglobin protein.

Dilution

IHC~~1:100~500

Format

Mouse IgG. Liquid in PBS containing 50% glycerol, 0.2% BSA and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-Myoglobin Antibody - Protein Information

Name MB (HGNC:6915)

Function

Monomeric heme protein which primary function is to store oxygen and facilitate its diffusion within muscle tissues. Reversibly binds oxygen through a pentacoordinated heme iron and enables its timely and efficient release as needed during periods of heightened demand (PubMed:30918256, PubMed:34679218). Depending on the oxidative conditions of tissues and cells, and in addition to its ability to bind oxygen, it also has a nitrite reductase activity whereby it regulates the production of bioactive nitric oxide (PubMed:32891753). Under stress conditions, like hypoxia and anoxia, it also protects cells against reactive oxygen species thanks to its pseudoperoxidase activity (PubMed:34679218).




Cellular LocationCytoplasm, sarcoplasm

Anti-Myoglobin Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Myoglobin Antibody - Images



Immunohistochemical analysis of Myoglobin staining in human cardiac muscle formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunohistochemical analysis of Myoglobin staining in human skeletal muscle formalin fixed





Tel: 858.875.1900 Fax: 858.875.1999

paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-Myoglobin Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within human Myoglobin. The exact sequence is proprietary.