

HSC70 Interacting Protein HIP (ST13) Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6247a

Specification

HSC70 Interacting Protein HIP (ST13) Antibody - Product Information

Application	WB, IHC-P,E
Primary Accession	P50502
Other Accession	Q8IZP2 , NP_003923
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	41332
Antigen Region	50-80

HSC70 Interacting Protein HIP (ST13) Antibody - Additional Information

Gene ID 6767

Other Names

Hsc70-interacting protein, Hip, Aging-associated protein 2, Progesterone receptor-associated p48 protein, Protein FAM10A1, Putative tumor suppressor ST13, Renal carcinoma antigen NY-REN-33, Suppression of tumorigenicity 13 protein, ST13, AAG2, FAM10A1, HIP, SNC6

Target/Specificity

This HSC70 Interacting Protein HIP (ST13) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 50-80 amino acids from human HSC70 Interacting Protein HIP (ST13).

Dilution

WB~~1:1000
IHC-P~~1:10~50
E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HSC70 Interacting Protein HIP (ST13) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

HSC70 Interacting Protein HIP (ST13) Antibody - Protein Information

Name ST13

Synonyms AAG2, FAM10A1, HIP, SNC6

Function One HIP oligomer binds the ATPase domains of at least two HSC70 molecules dependent on activation of the HSC70 ATPase by HSP40. Stabilizes the ADP state of HSC70 that has a high affinity for substrate protein. Through its own chaperone activity, it may contribute to the interaction of HSC70 with various target proteins (By similarity).

Cellular Location


Cytoplasm.

HSC70 Interacting Protein HIP (ST13) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HSC70 Interacting Protein HIP (ST13) Antibody - Images



Western blot analysis of anti-ST13 Antibody (N-term) (Cat.#AP6247a) pre-incubated with and without blocking peptide (BP6247a) in A2058 cell line lysate. ST13(arrow) was detected using the purified Pab .



Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with ST13 antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

HSC70 Interacting Protein HIP (ST13) Antibody - Background

ST13 is an adaptor protein that mediates the association of the heat shock proteins HSP70 and HSP90. This protein has been shown to be involved in the assembly process of glucocorticoid receptor, which requires the assistance of multiple molecular chaperones. The expression of this protein is reported to be downregulated in colorectal carcinoma tissue suggesting that is a candidate tumor suppressor.

HSC70 Interacting Protein HIP (ST13) Antibody - References

Rajapandi, T., et al., J. Biol. Chem. 275(29):22597-22604 (2000).
Morishima, Y., et al., J. Biol. Chem. 275(10):6894-6900 (2000).
Chen, S., et al., J. Biol. Chem. 273(52):35194-35200 (1998).
Johnson, B.D., et al., J. Biol. Chem. 273(6):3679-3686 (1998).
Cao, J., et al., J. Cancer Res. Clin. Oncol. 123(8):447-451 (1997).