

MEF2C Antibody (S387)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6285a

Specification

MEF2C Antibody (S387) - Product Information

Application	WB, IF, IHC-P,E
Primary Accession	Q06413
Other Accession	A4UTP7 , A0A096MJY4
Reactivity	Human, Mouse
Predicted	Pig, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	365-394

MEF2C Antibody (S387) - Additional Information

Gene ID 4208

Other Names

Myocyte-specific enhancer factor 2C, MEF2C

Target/Specificity

This MEF2C antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 365-394 amino acids from human MEF2C.

Dilution

WB~~1:1000-2000

IF~~1:10~50

IHC-P~~1:10~50

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MEF2C Antibody (S387) is for research use only and not for use in diagnostic or therapeutic procedures.

MEF2C Antibody (S387) - Protein Information

Name MEF2C ([HGNC:6996](#))

Function Transcription activator which binds specifically to the MEF2 element present in the regulatory regions of many muscle-specific genes. Controls cardiac morphogenesis and myogenesis, and is also involved in vascular development. Enhances transcriptional activation mediated by SOX18. Plays an essential role in hippocampal-dependent learning and memory by suppressing the number of excitatory synapses and thus regulating basal and evoked synaptic transmission. Crucial for normal neuronal development, distribution, and electrical activity in the neocortex. Necessary for proper development of megakaryocytes and platelets and for bone marrow B-lymphopoiesis. Required for B-cell survival and proliferation in response to BCR stimulation, efficient IgG1 antibody responses to T-cell-dependent antigens and for normal induction of germinal center B-cells. May also be involved in neurogenesis and in the development of cortical architecture (By similarity). Isoforms that lack the repressor domain are more active than isoform 1.

Cellular Location

Nucleus {ECO:0000250|UniProtKB:A0A096MJY4}. Cytoplasm, sarcoplasm {ECO:0000250|UniProtKB:A0A096MJY4}

Tissue Location


Expressed in brain and skeletal muscle.

MEF2C Antibody (S387) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MEF2C Antibody (S387) - Images




Fluorescent confocal image of HeLa cell stained with MEF2C Antibody (S387)(Cat#AP6285a).HeLa cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then

incubated with MEF2C primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C). Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7units/ml, 1 h at 37°C). Nuclei were counterstained with DAPI (blue) (10 µg/ml, 10 min). MEF2C immunoreactivity is localized to vesicles and Nucleus significantly.



All lanes : Anti-MEF2C-pS387. ctrl at 1:1000-2000 dilution Lane 1: K562 whole cell lysate Lane 2: RPMI 8226 whole cell lysate Lane 3: NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 51 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human brain tissue reacted with MEF2C Antibody (S387) (Cat.#AP6285a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

MEF2C Antibody (S387) - Background

MEF2C is a transcription activator which binds specifically to the MEF2 element present in the regulatory regions of many muscle-specific genes. This protein controls cardiac morphogenesis and myogenesis, and is also involved in vascular development. It may also be involved in neurogenesis and in the development of cortical architecture.

MEF2C Antibody (S387) - References

Konig, S., et al., *J. Biol. Chem.* 279(27):28187-28196 (2004). Maeda, T., et al., *J. Biol. Chem.* 277(50):48889-48898 (2002). Maeda, T., et al., *Biochem. Biophys. Res. Commun.* 294(4):791-797 (2002). Janson, C.G., et al., *Brain Res. Mol. Brain Res.* 97(1):70-82 (2001). Krainc, D., et al., *Genomics* 29(3):809-811 (1995).