

**Cleaved-MMP-14 (Y112) Polyclonal Antibody**  
**Catalog # AP63136****Specification****Cleaved-MMP-14 (Y112) Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P50281</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

**Cleaved-MMP-14 (Y112) Polyclonal Antibody - Additional Information****Gene ID** 4323**Other Names**

MMP14; Matrix metalloproteinase-14; MMP-14; MMP-X1; Membrane-type matrix metalloproteinase 1; MT-MMP 1; MTMMP1; Membrane-type-1 matrix metalloproteinase; MT1-MMP; MT1MMP

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**Cleaved-MMP-14 (Y112) Polyclonal Antibody - Protein Information****Name** MMP14**Function**

Endopeptidase that degrades various components of the extracellular matrix such as collagen. Activates progelatinase A. Essential for pericellular collagenolysis and modeling of skeletal and extraskeletal connective tissues during development (By similarity). May be involved in actin cytoskeleton reorganization by cleaving PTK7 (PubMed:<a href="http://www.uniprot.org/citations/20837484" target="\_blank">20837484</a>). Acts as a positive regulator of cell growth and migration via activation of MMP15. Involved in the formation of the fibrovascular tissues in association with pro-MMP2 (PubMed:<a href="http://www.uniprot.org/citations/12714657" target="\_blank">12714657</a>). Cleaves ADGRB1 to release vasculostatin-40 which inhibits angiogenesis (PubMed:<a href="http://www.uniprot.org/citations/22330140" target="\_blank">22330140</a>).

**Cellular Location**

Membrane; Single-pass type I membrane protein. Melanosome. Cytoplasm. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV Forms a complex with BST2 and localizes to the cytoplasm


**Tissue Location**

Expressed in stromal cells of colon, breast, and head and neck. Expressed in lung tumors.

**Cleaved-MMP-14 (Y112) Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Cleaved-MMP-14 (Y112) Polyclonal Antibody - Images****Cleaved-MMP-14 (Y112) Polyclonal Antibody - Background**

Endopeptidase that degrades various components of the extracellular matrix such as collagen. Activates progelatinase A. Essential for pericellular collagenolysis and modeling of skeletal and extracellular connective tissues during development (By similarity). May be involved in actin cytoskeleton reorganization by cleaving PTK7 (PubMed:20837484). Acts as a positive regulator of cell growth and migration via activation of MMP15. Involved in the formation of the fibrovascular tissues in association with pro-MMP2 (PubMed:12714657). Cleaves ADGRB1 to release vasculostatin-40 which inhibits angiogenesis (PubMed:22330140).