

SPPL3 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6313a**Specification**

SPPL3 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q8TCT6
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	50-81

SPPL3 Antibody (N-term) - Additional Information**Gene ID** 121665**Other Names**

Signal peptide peptidase-like 3, SPP-like 3, 3423-, Intramembrane protease 2, IMP-2, Presenilin homologous protein 1, PSH1, Presenilin-like protein 4, SPPL3, IMP2, PSL4

Target/Specificity

This SPPL3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 50-81 amino acids from the N-terminal region of human SPPL3.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SPPL3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SPPL3 Antibody (N-term) - Protein Information**Name** SPPL3 {ECO:0000303|PubMed:12077416, ECO:0000312|HGNC:HGNC:30424}

Function Intramembrane-cleaving aspartic protease (I-CLiP) that cleaves type II membrane protein substrates in or close to their luminal transmembrane domain boundaries (PubMed:[16873890](#), PubMed:[25354954](#), PubMed:[25827571](#)). Acts like a sheddase by mediating

the proteolytic release and secretion of active site-containing ectodomains of glycan-modifying glycosidase and glycosyltransferase enzymes such as MGAT5, B4GAT1 and B4GALT1 (PubMed:[25354954](#), PubMed:[25827571](#)). Catalyzes the intramembrane cleavage of the envelope glycoprotein gp130 and/or the leader peptide gp18LP of the simian foamy virus independent of prior ectodomain shedding by furin or furin-like proprotein convertase (PC)-mediated cleavage proteolysis (PubMed:[23132852](#)). May also have the ability to serve as a shedding protease for subsequent intramembrane proteolysis by SPPL2A and SPPL2B of the envelope glycoprotein gp130 (PubMed:[23132852](#)). Plays a role in the regulation of cellular glycosylation processes (PubMed:[25354954](#)). Required to link T-cell antigen receptor (TCR) and calcineurin-NFAT signaling cascades in lymphocytes by promoting the association of STIM1 and ORAI1 during store-operated calcium entry (SOCE) in a protease- independent manner (PubMed:[25384971](#)).

Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein. Golgi apparatus. Membrane; Multi-pass membrane protein; Luminal side

Tissue Location


Widely expressed (PubMed:15385547). Expressed in the brain (PubMed:11978763).

SPPL3 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SPPL3 Antibody (N-term) - Images



Western blot analysis of SPPL3 (arrow) using rabbit polyclonal SPPL3 Antibody (N-term) (Cat.#AP6313a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the SPPL3 gene (Lane 2) (Origene Technologies).

SPPL3 Antibody (N-term) - Background

Signal peptide peptidase (SPP) is an aspartyl protease that mediates clearance of signal peptides by proteolysis within the endoplasmic reticulum (ER). Like presenilins, SPP contains a critical GXGD motif in its C-terminal catalytic center. SPPL3 is one of several presenilin homologues/SPP-like proteins (PSHs/SPPL) that have been identified.

SPPL3 Antibody (N-term) - References

Grigorenko, A.P., et al., Biochemistry Mosc. 67(7):826-835 (2002).
Weihofen, A., et al., Science 296(5576):2215-2218 (2002).