

Cleaved-Factor Xa activated HC (I235) Polyclonal Antibody
Catalog # AP63158**Specification**

Cleaved-Factor Xa activated HC (I235) Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P00742
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Cleaved-Factor Xa activated HC (I235) Polyclonal Antibody - Additional Information**Gene ID** 2159**Other Names**

F10; Coagulation factor X; Stuart factor; Stuart-Prower factor

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Cleaved-Factor Xa activated HC (I235) Polyclonal Antibody - Protein Information**Name** F10**Function**

Factor Xa is a vitamin K-dependent glycoprotein that converts prothrombin to thrombin in the presence of factor Va, calcium and phospholipid during blood clotting.

Cellular Location

Secreted.

Tissue Location

Plasma; synthesized in the liver.



Cleaved-Factor Xa activated HC (I235) Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cleaved-Factor Xa activated HC (I235) Polyclonal Antibody - Images



Cleaved-Factor Xa activated HC (I235) Polyclonal Antibody - Background

Factor Xa is a vitamin K-dependent glycoprotein that converts prothrombin to thrombin in the presence of factor Va, calcium and phospholipid during blood clotting.