

**p73 (Acetyl Lys327) Polyclonal Antibody**  
Catalog # AP63235**Specification****p73 (Acetyl Lys327) Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">O15350</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

**p73 (Acetyl Lys327) Polyclonal Antibody - Additional Information****Gene ID** 7161**Other Names**

TP73; P73; Tumor protein p73; p53-like transcription factor; p53-related protein

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**p73 (Acetyl Lys327) Polyclonal Antibody - Protein Information****Name** TP73**Synonyms** P73**Function**

Participates in the apoptotic response to DNA damage. Isoforms containing the transactivation domain are pro-apoptotic, isoforms lacking the domain are anti-apoptotic and block the function of p53 and transactivating p73 isoforms. May be a tumor suppressor protein. Is an activator of FOXJ1 expression (By similarity). It is an essential factor for the positive regulation of lung ciliated cell differentiation (PubMed:<a href="http://www.uniprot.org/citations/34077761" target="\_blank">34077761</a>).

**Cellular Location**

Nucleus. Cytoplasm. Note=Accumulates in the nucleus in response to DNA damage

**Tissue Location**


Expressed in striatal neurons of patients with Huntington disease (at protein level). Brain, kidney, placenta, colon, heart, liver, spleen, skeletal muscle, prostate, thymus and pancreas Highly expressed in fetal tissue. Expressed in the respiratory epithelium (PubMed:34077761).

## p73 (Acetyl Lys327) Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## p73 (Acetyl Lys327) Polyclonal Antibody - Images



Western Blot analysis of Jurkat cells using Acetyl-p73 (K327) Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Western Blot analysis of JK cells using Acetyl-p73 (K327) Polyclonal Antibody. Secondary antibody was diluted at 1:20000

**p73 (Acetyl Lys327) Polyclonal Antibody - Background**

Participates in the apoptotic response to DNA damage. Isoforms containing the transactivation domain are pro-apoptotic, isoforms lacking the domain are anti-apoptotic and block the function of p53 and transactivating p73 isoforms. May be a tumor suppressor protein.