

# Cytokeratin 18 Monoclonal Antibody(2F7)

**Catalog # AP63350** 

### **Specification**

# Cytokeratin 18 Monoclonal Antibody(2F7) - Product Information

Application WB
Primary Accession P05783

Reactivity Human, Mouse

Host Mouse Clonality Monoclonal

## Cytokeratin 18 Monoclonal Antibody(2F7) - Additional Information

**Gene ID 3875** 

**Other Names** 

KRT18; CYK18; PIG46; Keratin, type I cytoskeletal 18; Cell proliferation-inducing gene 46 protein;

Cytokeratin-18; CK-18; Keratin-18; K18

**Dilution** 

WB~~WB: 1:500-1000

**Format** 

PBS, pH 7.4, containing 0.09% (W/V) sodium azide as Preservative and 50% Glycerol.

**Storage Conditions** 

-20°C

### Cytokeratin 18 Monoclonal Antibody(2F7) - Protein Information

Name KRT18

**Synonyms** CYK18

#### **Function**

Involved in the uptake of thrombin-antithrombin complexes by hepatic cells (By similarity). When phosphorylated, plays a role in filament reorganization. Involved in the delivery of mutated CFTR to the plasma membrane. Together with KRT8, is involved in interleukin-6 (IL-6)-mediated barrier protection.

### **Cellular Location**

Nucleus matrix {ECO:0000250|UniProtKB:Q5BJY9}. Cytoplasm, perinuclear region. Nucleus, nucleolus. Cytoplasm {ECO:0000250|UniProtKB:Q5BJY9}

## **Tissue Location**

Expressed in colon, placenta, liver and very weakly in exocervix. Increased expression observed in lymph nodes of breast carcinoma.




## Cytokeratin 18 Monoclonal Antibody(2F7) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### Cytokeratin 18 Monoclonal Antibody(2F7) - Images



Cytokeratin 18 Monoclonal Antibody(2F7) - Background

Involved in the uptake of thrombin-antithrombin complexes by hepatic cells (By similarity). When phosphorylated, plays a role in filament reorganization. Involved in the delivery of mutated CFTR to the plasma membrane. Together with KRT8, is involved in interleukin-6 (IL-6)-mediated barrier protection.