

eNOS Monoclonal Antibody(Mix)
Catalog # AP63385**Specification****eNOS Monoclonal Antibody(Mix) - Product Information**

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | P29474 |
| Reactivity | Human, Mouse, Rat |
| Host | Mouse |
| Clonality | Monoclonal |

eNOS Monoclonal Antibody(Mix) - Additional Information**Gene ID 4846****Other Names**

NOS3; Nitric oxide synthase, endothelial; Constitutive NOS; cNOS; EC-NOS; Endothelial NOS; eNOS; NOS type III; NOSIII

Dilution

WB~~WB: 1:500-2000

Format

PBS, pH 7.4, containing 0.09% (W/V) sodium azide as Preservative and 50% Glycerol.

Storage Conditions

-20°C

eNOS Monoclonal Antibody(Mix) - Protein Information**Name NOS3 ([HGNC:7876](#))****Function**

Produces nitric oxide (NO) which is implicated in vascular smooth muscle relaxation through a cGMP-mediated signal transduction pathway (PubMed:1378832). NO mediates vascular endothelial growth factor (VEGF)-induced angiogenesis in coronary vessels and promotes blood clotting through the activation of platelets.

Cellular Location

Cell membrane. Membrane, caveola. Cytoplasm, cytoskeleton. Golgi apparatus. Note=Specifically associates with actin cytoskeleton in the G2 phase of the cell cycle; which is favored by interaction with NOSIP and results in a reduced enzymatic activity

Tissue Location


Platelets, placenta, liver and kidney.

eNOS Monoclonal Antibody(Mix) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

eNOS Monoclonal Antibody(Mix) - Images



Western blot analysis of Rat Heart Tissue, diluted at 1:1000.



The picture was kindly provided by our customer

eNOS Monoclonal Antibody(Mix) - Background

Produces nitric oxide (NO) which is implicated in vascular smooth muscle relaxation through a cGMP-mediated signal transduction pathway. NO mediates vascular endothelial growth factor (VEGF)-induced angiogenesis in coronary vessels and promotes blood clotting through the activation of platelets.