

EPG5 Polyclonal Antibody
Catalog # AP63518**Specification**

EPG5 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q9HCE0
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

EPG5 Polyclonal Antibody - Additional Information**Gene ID** 57724**Other Names**

EPG5; KIAA1632; Ectopic P granules protein 5 homolog

Dilution

WB~~WB: 1:1000-2000 IHC: 1:200-500

IHC-P~~N/A

Format

PBS, pH 7.4, containing 0.09% (W/V) sodium azide as Preservative and 50% Glycerol.

Storage Conditions

-20°C

EPG5 Polyclonal Antibody - Protein Information**Name** EPG5**Synonyms** KIAA1632**Function**

Involved in autophagy. May play a role in a late step of autophagy, such as clearance of autophagosomal cargo. Plays a key role in innate and adaptive immune response triggered by unmethylated cytidine-phosphate-guanosine (CpG) dinucleotides from pathogens, and mediated by the nucleotide-sensing receptor TLR9. It is necessary for the translocation of CpG dinucleotides from early endosomes to late endosomes and lysosomes, where TLR9 is located (PubMed:29130391).

Cellular Location



Cytoplasm, perinuclear region. Lysosome

EPG5 Polyclonal Antibody - Protocols



Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EPG5 Polyclonal Antibody - Images



170KD —
130KD —
95KD —
70KD —



EPG5 Polyclonal Antibody - Background

Involved in autophagy. May play a role in a late step of autophagy, such as clearance of autophagosomal cargo.