

Annexin V Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6580a**Specification**

Annexin V Antibody (N-term) - Product Information

Application	IHC-P-Leica, FC, WB,E
Primary Accession	P08758
Other Accession	Q4R4H7 , P81287
Reactivity	Human, Mouse
Predicted	Bovine, Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	1-30

Annexin V Antibody (N-term) - Additional Information**Gene ID** 308**Other Names**

Annexin A5, Anchorin CII, Annexin V, Annexin-5, Calphobindin I, CBP-I, Endonexin II, Lipocortin V, Placental anticoagulant protein 4, PP4, Placental anticoagulant protein I, PAP-I, Thromboplastin inhibitor, Vascular anticoagulant-alpha, VAC-alpha, ANXA5, ANX5, ENX2, PP4

Target/Specificity

This Annexin V antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human Annexin V.

Dilution

IHC-P-Leica~~1:500

FC~~1:10~50

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Annexin V Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Annexin V Antibody (N-term) - Protein Information

Name ANXA5

Synonyms ANX5, ENX2, PP4


Function This protein is an anticoagulant protein that acts as an indirect inhibitor of the thromboplastin-specific complex, which is involved in the blood coagulation cascade.

Annexin V Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Annexin V Antibody (N-term) - Images




All lanes : Anti-Annexin V Antibody (N-term) at 1:2000 dilution Lane 1: HepG2 whole cell lysate Lane 2: 293 whole cell lysate Lane 3: U-2OS whole cell lysate Lane 4: A2058 whole cell lysate Lane 5: T47D whole cell lysate Lane 6: Mouse lung lysate Lane 7: Mouse heart lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 36 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of paraffin-embedded Human hepatocarcinoma tissue using AP6580A performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunohistochemical analysis of paraffin-embedded Human placenta tissue using AP6580A performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Annexin V Antibody (N-term) (Cat. #AP6580a) flow cytometric analysis of A2058 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Annexin V Antibody (N-term) - Background

Annexin 5 belongs to the annexin family of calcium-dependent phospholipid binding proteins some of which have been implicated in membrane-related events along exocytotic and endocytotic pathways. Annexin 5 is a phospholipase A2 and protein kinase C inhibitory protein with calcium channel activity and a potential role in cellular signal transduction, inflammation, growth and differentiation. Annexin 5 has also been described as placental anticoagulant protein I, vascular anticoagulant-alpha, endonexin II, lipocortin V, placental protein 4 and anchorin CII.

Annexin V Antibody (N-term) - References

Tiscia, G., Thromb. Haemost. 102 (2), 309-313 (2009)
Kim, T.H., Bone 45 (1), 125-131 (2009)