

MCT-1 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6655a**Specification**

MCT-1 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q9ULC4
Other Accession	Q9DB27 , Q7ZV34 , Q5ZI42 , Q2KIE4 , Q5PPY1
Reactivity	Human, Mouse
Predicted	Xenopus, Bovine, Chicken, Zebrafish
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	20555
Antigen Region	1-30

MCT-1 Antibody (N-term) - Additional Information**Gene ID** 28985**Other Names**

Malignant T-cell-amplified sequence 1, MCT-1, Multiple copies T-cell malignancies, MCTS1, MCT1

Target/Specificity

This MCT-1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human MCT-1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MCT-1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MCT-1 Antibody (N-term) - Protein Information**Name** MCTS1**Synonyms** MCT1

Function Anti-oncogene that plays a role in cell cycle regulation; decreases cell doubling time and anchorage-dependent growth; shortens the duration of G1 transit time and G1/S transition. When constitutively expressed, increases CDK4 and CDK6 kinases activity and CCND1/cyclin D1 protein level, as well as G1 cyclin/CDK complex formation. Involved in translation initiation; promotes recruitment of aminoacylated initiator tRNA to P site of 40S ribosomes. Can promote release of deacylated tRNA and mRNA from recycled 40S subunits following ABCE1-mediated dissociation of post-termination ribosomal complexes into subunits. Plays a role as translation enhancer; recruits the density-regulated protein/DENR and binds to the cap complex of the 5'-terminus of mRNAs, subsequently altering the mRNA translation profile; up-regulates protein levels of BCL2L2, TFDP1, MRE11, CCND1 and E2F1, while mRNA levels remains constant. Hyperactivates DNA damage signaling pathway; increased gamma-irradiation-induced phosphorylation of histone H2AX, and induces damage foci formation. Increases the overall number of chromosomal abnormalities such as larger chromosomes formation and multiple chromosomal fusions when overexpressed in gamma- irradiated cells. May play a role in promoting lymphoid tumor development: lymphoid cell lines overexpressing MCTS1 exhibit increased growth rates and display increased protection against apoptosis. May contribute to the pathogenesis and progression of breast cancer via promotion of angiogenesis through the decline of inhibitory THBS1/thrombospondin-1, and inhibition of apoptosis. Involved in the process of proteasome degradation to down-regulate Tumor suppressor p53/TP53 in breast cancer cell; Positively regulates phosphorylation of MAPK1 and MAPK3. Involved in translation initiation; promotes aminoacylated initiator tRNA to P site of 40S ribosomes. Can promote release of deacylated tRNA and mRNA from recycled 40S subunits following ABCE1-mediated dissociation of post-termination ribosomal complexes into subunits.

Cellular Location

Cytoplasm. Note=Nuclear relocalization after DNA damage

Tissue Location


Ubiquitous. Over-expressed in T-cell lymphoid cell lines and in non-Hodgkin lymphoma cell lines as well as in a subset of primary large B-cell lymphomas.

MCT-1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MCT-1 Antibody (N-term) - Images



Western blot analysis of MCT-1 antibody (N-term) (Cat. #AP6655a) in mouse bladder tissue lysates (35ug/lane). MCTS1 (arrow) was detected using the purified Pab.

MCT-1 Antibody (N-term) - Background

MCTS1 play a role in cell cycle regulation; decreases cell doubling time and anchorage-dependent growth; shortens the duration of G1 transit time and G1/S transition.

MCT-1 Antibody (N-term) - References

Kasiappan,R., Mol. Cancer Res. 7 (4), 536-548 (2009)
Mazan-Mamczarz,K., Leuk. Res. 33 (3), 474-482 (2009)
Shi,B., J. Cell. Biochem. 90 (1), 68-79 (2003)