

### **KRT10 Antibody (Center)**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP6704c

### Specification

# KRT10 Antibody (Center) - Product Information

Application	FC, IHC-P, WB,E
Primary Accession	<u>P13645</u>
Other Accession	<u>Q6IFW6</u>
Reactivity	Human
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	58827
Antigen Region	318-346

## KRT10 Antibody (Center) - Additional Information

Gene ID 3858

Other Names Keratin, type I cytoskeletal 10, Cytokeratin-10, CK-10, Keratin-10, K10, KRT10, KPP

Target/Specificity

This KRT10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 318-346 amino acids from the Central region of human KRT10.

Dilution FC~~1:10~50 IHC-P~~1:50~100 WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

KRT10 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

# KRT10 Antibody (Center) - Protein Information



## Name KRT10

### Synonyms KPP

**Function** Plays a role in the establishment of the epidermal barrier on plantar skin (By similarity). Involved in the maintenance of cell layer development and keratin filament bundles in suprabasal cells of the epithelium (By similarity).

**Cellular Location** Secreted, extracellular space. Cell surface. Cytoplasm

#### **Tissue Location**

Seen in all suprabasal cell layers including stratum corneum. Expressed on the surface of lung cell lines (PubMed:19627498). Localized on the surface of desquamated nasal epithelial cells (at protein level) (PubMed:12427098)

## **KRT10 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

## KRT10 Antibody (Center) - Images



Western blot analysis of KRT10 antibody (Center) (Cat. #AP6704c) in CEM cell line lysates (35ug/lane). KRT10 (arrow) was detected using the purified Pab.





Formalin-fixed and paraffin-embedded human skin reacted with KRT10 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



KRT10 Antibody (Center) (Cat. #AP6704c) flow cytometric analysis of CEM cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

# KRT10 Antibody (Center) - Background

KRT10 is a member of the type I (acidic) cytokeratin family, which belongs to the superfamily of intermediate filament (IF) proteins. Keratins are heteropolymeric structural proteins which form the intermediate filament. These filaments, along with actin microfilaments and microtubules, compose the cytoskeleton of epithelial cells. Mutations in its gene are associated with epidermolytic hyperkeratosis.

# **KRT10 Antibody (Center) - References**

Morais,P., Eur J Dermatol 19 (4), 333-336 (2009) Barcelos,A.C., J. Cutan. Pathol. 36 (6), 647-654 (2009) **KRT10 Antibody (Center) - Citations** 

• <u>Differential miRNA expression profiles in human keratinocytes in response to protein kinase</u> <u>C inhibitor.</u>