

MVD Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6717a**Specification**

MVD Antibody (N-term) - Product Information

Application	IHC-P, WB,E
Primary Accession	P53602
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	43405
Antigen Region	7-34

MVD Antibody (N-term) - Additional Information**Gene ID** 4597**Other Names**

Diphosphomevalonate decarboxylase, Mevalonate (diphospho)decarboxylase, MDDase, Mevalonate pyrophosphate decarboxylase, MVD, MPD

Target/Specificity

This MVD antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 7-34 amino acids from the N-terminal region of human MVD.

Dilution

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MVD Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MVD Antibody (N-term) - Protein Information**Name** MVD

Synonyms MPD {ECO:0000303|PubMed:14972328}

Function Catalyzes the ATP dependent decarboxylation of (R)-5- diphosphomevalonate to form isopentenyl diphosphate (IPP). Functions in the mevalonate (MVA) pathway leading to isopentenyl diphosphate (IPP), a key precursor for the biosynthesis of isoprenoids and sterol synthesis.

Cellular Location

Cytoplasm.

Tissue Location


Expressed in heart, skeletal muscle, lung, liver, brain, pancreas, kidney and placenta.

MVD Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MVD Antibody (N-term) - Images



Western blot analysis of MVD (arrow) using rabbit polyclonal MVD Antibody (N-term) (Cat. #AP6717a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the MVD gene (Lane 2).



Formalin-fixed and paraffin-embedded human Skeletal muscle reacted with MVD Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

MVD Antibody (N-term) - Background

The enzyme mevalonate pyrophosphate decarboxylase catalyzes the conversion of mevalonate pyrophosphate into isopentenyl pyrophosphate in one of the early steps in cholesterol biosynthesis. It decarboxylates and dehydrates its substrate while hydrolyzing ATP.

MVD Antibody (N-term) - References

Voynova,N.E., Arch. Biochem. Biophys. 480 (1), 58-67 (2008)
Hogenboom,S., Mol. Genet. Metab. 81 (3), 216-224 (2004)
Wadhwa,R., Biochem. Biophys. Res. Commun. 302 (4), 735-742 (2003)