

**Btk (phospho Tyr551) Polyclonal Antibody**  
**Catalog # AP67220****Specification**

---

**Btk (phospho Tyr551) Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q06187</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

**Btk (phospho Tyr551) Polyclonal Antibody - Additional Information****Gene ID** 695**Other Names**

BTK; AGMX1; ATK; BPK; Tyrosine-protein kinase BTK; Agammaglobulinaemia tyrosine kinase; ATK; B-cell progenitor kinase; BPK; Bruton tyrosine kinase

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**Btk (phospho Tyr551) Polyclonal Antibody - Protein Information****Name** BTK**Synonyms** AGMX1, ATK, BPK**Function**

Non-receptor tyrosine kinase indispensable for B lymphocyte development, differentiation and signaling (PubMed:<a href="http://www.uniprot.org/citations/19290921" target="\_blank">19290921</a>). Binding of antigen to the B-cell antigen receptor (BCR) triggers signaling that ultimately leads to B-cell activation (PubMed:<a href="http://www.uniprot.org/citations/19290921" target="\_blank">19290921</a>). After BCR engagement and activation at the plasma membrane, phosphorylates PLCG2 at several sites, igniting the downstream signaling pathway through calcium mobilization, followed by activation of the protein kinase C (PKC) family members (PubMed:<a href="http://www.uniprot.org/citations/11606584" target="\_blank">11606584</a>). PLCG2 phosphorylation is performed in close cooperation with the adapter protein B-cell linker protein BLNK (PubMed:<a href="http://www.uniprot.org/citations/11606584" target="\_blank">11606584</a>). BTK acts as a platform to bring together a diverse array of signaling proteins and is implicated in cytokine receptor signaling pathways (PubMed:<a href="http://www.uniprot.org/citations/11606584" target="\_blank">11606584</a>).

[16517732](http://www.uniprot.org/citations/16517732), PubMed: [17932028](http://www.uniprot.org/citations/17932028)). Plays an important role in the function of immune cells of innate as well as adaptive immunity, as a component of the Toll-like receptors (TLR) pathway (PubMed: [16517732](http://www.uniprot.org/citations/16517732)). The TLR pathway acts as a primary surveillance system for the detection of pathogens and are crucial to the activation of host defense (PubMed: [16517732](http://www.uniprot.org/citations/16517732)). Especially, is a critical molecule in regulating TLR9 activation in splenic B-cells (PubMed: [16517732](http://www.uniprot.org/citations/16517732), PubMed: [17932028](http://www.uniprot.org/citations/17932028)). Within the TLR pathway, induces tyrosine phosphorylation of TIRAP which leads to TIRAP degradation (PubMed: [16415872](http://www.uniprot.org/citations/16415872)). BTK also plays a critical role in transcription regulation (PubMed: [19290921](http://www.uniprot.org/citations/19290921)). Induces the activity of NF- $\kappa$ B, which is involved in regulating the expression of hundreds of genes (PubMed: [19290921](http://www.uniprot.org/citations/19290921)). BTK is involved on the signaling pathway linking TLR8 and TLR9 to NF- $\kappa$ B (PubMed: [19290921](http://www.uniprot.org/citations/19290921)). Acts as an activator of NLRP3 inflammasome assembly by mediating phosphorylation of NLRP3 (PubMed: [34554188](http://www.uniprot.org/citations/34554188)). Transiently phosphorylates transcription factor GTF2I on tyrosine residues in response to BCR (PubMed: [9012831](http://www.uniprot.org/citations/9012831)). GTF2I then translocates to the nucleus to bind regulatory enhancer elements to modulate gene expression (PubMed: [9012831](http://www.uniprot.org/citations/9012831)). ARID3A and NFAT are other transcriptional target of BTK (PubMed: [16738337](http://www.uniprot.org/citations/16738337)). BTK is required for the formation of functional ARID3A DNA-binding complexes (PubMed: [16738337](http://www.uniprot.org/citations/16738337)). There is however no evidence that BTK itself binds directly to DNA (PubMed: [16738337](http://www.uniprot.org/citations/16738337)). BTK has a dual role in the regulation of apoptosis (PubMed: [9751072](http://www.uniprot.org/citations/9751072)). Plays a role in STING1- mediated induction of type I interferon (IFN) response by phosphorylating DDX41 (PubMed: [25704810](http://www.uniprot.org/citations/25704810)).

### Cellular Location

Cytoplasm. Cell membrane; Peripheral membrane protein. Nucleus Membrane raft {ECO:0000250|UniProtKB:P35991}. Note=In steady state, BTK is predominantly cytosolic. Following B-cell receptor (BCR) engagement by antigen, translocates to the plasma membrane through its PH domain Plasma membrane localization is a critical step in the activation of BTK. A fraction of BTK also shuttles between the nucleus and the cytoplasm, and nuclear export is mediated by the nuclear export receptor CRM1.

### Tissue Location

Predominantly expressed in B-lymphocytes.


### Btk (phospho Tyr551) Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Btk (phospho Tyr551) Polyclonal Antibody - Images**



### **Btk (phospho Tyr551) Polyclonal Antibody - Background**

Non-receptor tyrosine kinase indispensable for B lymphocyte development, differentiation and signaling. Binding of antigen to the B-cell antigen receptor (BCR) triggers signaling that ultimately leads to B-cell activation. After BCR engagement and activation at the plasma membrane, phosphorylates PLCG2 at several sites, igniting the downstream signaling pathway through calcium mobilization, followed by activation of the protein kinase C (PKC) family members. PLCG2 phosphorylation is performed in close cooperation with the adapter protein B-cell linker protein BLNK. BTK acts as a platform to bring together a diverse array of signaling proteins and is implicated in cytokine receptor signaling pathways. Plays an important role in the function of immune cells of innate as well as adaptive immunity, as a component of the Toll-like receptors (TLR) pathway. The TLR pathway acts as a primary surveillance system for the detection of pathogens and are crucial to the activation of host defense. Especially, is a critical molecule in regulating TLR9 activation in splenic B-cells. Within the TLR pathway, induces tyrosine phosphorylation of TIRAP which leads to TIRAP degradation. BTK plays also a critical role in transcription regulation. Induces the activity of NF-kappa-B, which is involved in regulating the expression of hundreds of genes. BTK is involved on the signaling pathway linking TLR8 and TLR9 to NF-kappa-B. Transiently phosphorylates transcription factor GTF2I on tyrosine residues in response to BCR. GTF2I then translocates to the nucleus to bind regulatory enhancer elements to modulate gene expression. ARID3A and NFAT are other transcriptional target of BTK. BTK is required for the formation of functional ARID3A DNA-binding complexes. There is however no evidence that BTK itself binds directly to DNA. BTK has a dual role in the regulation of apoptosis.