

KIF20A (phospho Ser528) Polyclonal Antibody
Catalog # AP67305**Specification**

KIF20A (phospho Ser528) Polyclonal Antibody - Product Information

| | |
|-------------------|------------------------|
| Application | WB, IHC-P |
| Primary Accession | O95235 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |

KIF20A (phospho Ser528) Polyclonal Antibody - Additional Information**Gene ID** 10112**Other Names**

KIF20A; MKLP2; RAB6KIFL; Kinesin-like protein KIF20A; GG10_2; Mitotic kinesin-like protein 2; MKlp2; Rab6-interacting kinesin-like protein; Rabkinesin-6

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

KIF20A (phospho Ser528) Polyclonal Antibody - Protein Information**Name** KIF20A**Synonyms** MKLP2, RAB6KIFL**Function**

Mitotic kinesin required for chromosome passenger complex (CPC)-mediated cytokinesis. Following phosphorylation by PLK1, involved in recruitment of PLK1 to the central spindle. Interacts with guanosine triphosphate (GTP)-bound forms of RAB6A and RAB6B. May act as a motor required for the retrograde RAB6 regulated transport of Golgi membranes and associated vesicles along microtubules. Has a microtubule plus end- directed motility.

Cellular Location



Golgi apparatus. Cytoplasm, cytoskeleton, spindle. Note=Localizes to the spindle midzone during anaphase and telophase.

KIF20A (phospho Ser528) Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KIF20A (phospho Ser528) Polyclonal Antibody - Images



KIF20A (phospho Ser528) Polyclonal Antibody - Background

Mitotic kinesin required for chromosome passenger complex (CPC)-mediated cytokinesis.

Following phosphorylation by PLK1, involved in recruitment of PLK1 to the central spindle. Interacts with guanosine triphosphate (GTP)-bound forms of RAB6A and RAB6B. May act as a motor required for the retrograde RAB6 regulated transport of Golgi membranes and associated vesicles along microtubules. Has a microtubule plus end-directed motility.