

**KIR6.2 (phospho Thr224) Polyclonal Antibody**  
**Catalog # AP67494****Specification****KIR6.2 (phospho Thr224) Polyclonal Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC-P, IF          |
| Primary Accession | <a href="#">Q14654</a> |
| Reactivity        | Human, Mouse, Rat      |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |

**KIR6.2 (phospho Thr224) Polyclonal Antibody - Additional Information****Gene ID** 3767**Other Names**

KCNJ11; ATP-sensitive inward rectifier potassium channel 11; IKATP; Inward rectifier K(+) channel Kir6.2; Potassium channel; inwardly rectifying subfamily J member 11

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.  
IHC-P~~N/A  
IF~~1:50~200

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**KIR6.2 (phospho Thr224) Polyclonal Antibody - Protein Information****Name** KCNJ11**Function**

Inward rectifier potassium channel that forms the pore of ATP-sensitive potassium channels (KATP), regulating potassium permeability as a function of cytoplasmic ATP and ADP concentrations in many different cells (PubMed:<a href="http://www.uniprot.org/citations/29286281" target="\_blank">29286281</a>, PubMed:<a href="http://www.uniprot.org/citations/34815345" target="\_blank">34815345</a>). Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium; as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium. Can be blocked by extracellular barium (By similarity). In pancreatic cells, it forms KATP channels with ABCC8/SUR1 (PubMed:<a href="http://www.uniprot.org/citations/29286281" target="\_blank">29286281</a>, PubMed:<a href="http://www.uniprot.org/citations/29286281" target="\_blank">29286281</a>, PubMed:<a href="http://www.uniprot.org/citations/29286281" target="\_blank">29286281</a>).

href="http://www.uniprot.org/citations/34815345" target="\_blank">34815345</a>). Can form cardiac and smooth muscle-type KATP channels with ABCC9.

#### Cellular Location


Membrane; Multi-pass membrane protein.

### KIR6.2 (phospho Thr224) Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### KIR6.2 (phospho Thr224) Polyclonal Antibody - Images



### KIR6.2 (phospho Thr224) Polyclonal Antibody - Background

This receptor is controlled by G proteins. Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium; as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium. Can be blocked by extracellular barium (By similarity). Subunit of ATP-sensitive potassium channels (KATP). Can form cardiac and smooth muscle-type KATP channels with ABCC9. KCNJ11 forms the channel pore while ABCC9 is required for activation and regulation.