

Integrin β 4 (phospho Tyr1510) Polyclonal Antibody
Catalog # AP67687**Specification****Integrin β 4 (phospho Tyr1510) Polyclonal Antibody - Product Information**

Application	WB, IHC-P
Primary Accession	P16144
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Integrin β 4 (phospho Tyr1510) Polyclonal Antibody - Additional Information**Gene ID** 3691**Other Names**

ITGB4; Integrin beta-4; GP150; CD antigen CD104

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Integrin β 4 (phospho Tyr1510) Polyclonal Antibody - Protein Information**Name** ITGB4**Function**

Integrin alpha-6/beta-4 is a receptor for laminin. Plays a critical structural role in the hemidesmosome of epithelial cells. Is required for the regulation of keratinocyte polarity and motility. ITGA6:ITGB4 binds to NRG1 (via EGF domain) and this binding is essential for NRG1-ERBB signaling (PubMed:<[a href="http://www.uniprot.org/citations/20682778" target="_blank">20682778](http://www.uniprot.org/citations/20682778)). ITGA6:ITGB4 binds to IGF1 and this binding is essential for IGF1 signaling (PubMed:<[a href="http://www.uniprot.org/citations/22351760" target="_blank">22351760](http://www.uniprot.org/citations/22351760)). ITGA6:ITGB4 binds to IGF2 and this binding is essential for IGF2 signaling (PubMed:<[a href="http://www.uniprot.org/citations/28873464" target="_blank">28873464](http://www.uniprot.org/citations/28873464)).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell membrane; Lipid-anchor. Cell junction, hemidesmosome Note=Colocalizes with DST at the leading edge of migrating keratinocytes


Tissue Location


Integrin alpha-6/beta-4 is predominantly expressed by epithelia. Isoform beta-4D is also expressed in colon and placenta Isoform beta-4E is also expressed in epidermis, lung, duodenum, heart, spleen and stomach

Integrin β 4 (phospho Tyr1510) Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Integrin β 4 (phospho Tyr1510) Polyclonal Antibody - Images



Integrin β 4 (phospho Tyr1510) Polyclonal Antibody - Background

Integrin alpha-6/beta-4 is a receptor for laminin. Plays a critical structural role in the hemidesmosome of epithelial cells. Is required for the regulation of keratinocyte polarity and motility. ITGA6:ITGB4 binds to NRG1 (via EGF domain) and this binding is essential for NRG1-ERBB signaling (PubMed:20682778). ITGA6:ITGB4 binds to IGF1 and this binding is essential for IGF1 signaling (PubMed:22351760). ITGA6:ITGB4 binds to IGF2 and this binding is essential for IGF2 signaling (PubMed:28873464).