

PFTK1 Antibody (N-term P82) Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP6780a

Specification

PFTK1 Antibody (N-term P82) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype	WB, FC, IHC-P,E <u>094921</u> <u>035495</u> Human Mouse Rabbit Polyclonal Rabbit IgG
•	
	_
Antigen Region	67-96

PFTK1 Antibody (N-term P82) - Additional Information

Gene ID 5218

Other Names

Cyclin-dependent kinase 14, Cell division protein kinase 14, Serine/threonine-protein kinase PFTAIRE-1, hPFTAIRE1, CDK14, KIAA0834, PFTK1

Target/Specificity

This PFTK1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 67-96 amino acids from the N-terminal region of human PFTK1.

Dilution WB~~1:1000 FC~~1:10~50 IHC-P~~1:100 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PFTK1 Antibody (N-term P82) is for research use only and not for use in diagnostic or therapeutic procedures.

PFTK1 Antibody (N-term P82) - Protein Information



Name CDK14

Synonyms KIAA0834, PFTK1

Function Serine/threonine-protein kinase involved in the control of the eukaryotic cell cycle, whose activity is controlled by an associated cyclin. Acts as a cell-cycle regulator of Wnt signaling pathway during G2/M phase by mediating the phosphorylation of LRP6 at 'Ser-1490', leading to the activation of the Wnt signaling pathway. Acts as a regulator of cell cycle progression and cell proliferation via its interaction with CCDN3. Phosphorylates RB1 in vitro, however the relevance of such result remains to be confirmed in vivo. May also play a role in meiosis, neuron differentiation and may indirectly act as a negative regulator of insulin-responsive glucose transport.

Cellular Location

Cell membrane; Peripheral membrane protein. Cytoplasm. Nucleus. Note=Recruited to the cell membrane by CCNY

Tissue Location


Highly expressed in brain, pancreas, kidney, heart, testis and ovary. Also detected at lower levels in other tissues except in spleen and thymus where expression is barely detected

PFTK1 Antibody (N-term P82) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>


PFTK1 Antibody (N-term P82) - Images




Western blot analysis of lysate from human brain tissue lysate, using PFTK1 Antibody (N-term P82)(Cat. #AP6780a). AP6780a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG




H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug per lane.



PFTK1 Antibody (N-term P82) (RB18842) IHC analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the PFTK1 Antibody (N-term P82) for immunohistochemistry. Clinical relevance has not been evaluated.




Formalin-fixed and paraffin-embedded H.brain tissue reacted with PFTK1 Antibody (N-term P82) (Cat#AP6780a).



Formalin-fixed and paraffin-embedded H.testis tissue reacted with PFTK1 Antibody (N-term P82) (Cat#AP6780a).





PFTK1 Antibody (N-term P82) (Cat. #AP6780a) flow cytometric analysis of HepG2 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

PFTK1 Antibody (N-term P82) - Background

PFTK1 is a member of the CDC2 (MIM 116940)-related protein kinase family. It May play a role in meiosis as well as in neuron differentiation and/or function (By similarity).

PFTK1 Antibody (N-term P82) - References

Denoeud, F., et.al., Genome Res. 17 (6), 746-759 (2007)