

TPTEa Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6811a**Specification**

TPTEa Antibody (N-term) - Product Information

Application	IHC-P, WB,E
Primary Accession	P56180
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	64322
Antigen Region	42-74

TPTEa Antibody (N-term) - Additional Information**Gene ID** 7179**Other Names**

Putative tyrosine-protein phosphatase TPTE, Cancer/testis antigen 44, CT44, Transmembrane phosphatase with tensin homology, Tumor antigen BJ-HCC-5, TPTE

Target/Specificity

This TPTEa antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 42-74 amino acids from the N-terminal region of human TPTEa.

Dilution

IHC-P~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TPTEa Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TPTEa Antibody (N-term) - Protein Information**Name** TPTE

Function Could be involved in signal transduction.

Cellular Location

Membrane; Multi-pass membrane protein.

Tissue Location


Exclusively expressed in testis.

TPTEa Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TPTEa Antibody (N-term) - Images



TPTEa Antibody (E57) (Cat. #AP6811a) western blot analysis in MDA-MB231 cell line lysates (35ug/lane). This demonstrates the TPTEa antibody detected the TPTEa protein (arrow).



TPTEa Antibody (N-term) (Cat. #AP6811a) immunohistochemistry analysis in formalin fixed and paraffin embedded human testis tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of TPTEa Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

TPTEa Antibody (N-term) - Background

Ubiquitin is a 76 amino acid highly conserved eukaryotic polypeptide that selectively marks cellular proteins for proteolytic degradation by the 26S proteasome. The process of target selection, covalent attachment and shuttle to the 26S proteasome is a vital means of regulating the concentrations of key regulatory proteins in the cell by limiting their lifespans. Polyubiquitination is a common feature of this modification. Serial steps for modification include the activation of ubiquitin, an ATP-dependent formation of a thioester bond between ubiquitin and the enzyme E1, transfer by transacylation of ubiquitin from E1 to the ubiquitin conjugating enzyme E2, and covalent linkage to the target protein directly by E2 or via E3 ligase enzyme. Deubiquitination enzymes also exist to reverse the marking of protein substrates. Posttranslational tagging by Ub is involved in a multitude of cellular processes, including the cell cycle, cell growth and differentiation, embryogenesis, apoptosis, signal transduction, DNA repair, regulation of transcription and DNA replication, transmembrane transport, stress responses, the immune response, and nervous system functions.

TPTEa Antibody (N-term) - References

Hattori, M., et al., Nature 405(6784):311-319 (2000).
Chen, H., et al., Hum. Genet. 105(5):399-409 (1999).