

Nuclear Receptor NR4A1 (Phospho-Ser351) Antibody
Catalog # AP68180**Specification****Nuclear Receptor NR4A1 (Phospho-Ser351) Antibody - Product Information**

Application	WB
Primary Accession	P22736
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Nuclear Receptor NR4A1 (Phospho-Ser351) Antibody - Additional Information**Gene ID** 3164**Other Names**

Nuclear receptor subfamily 4 group A member 1 (Early response protein NAK1) (Nuclear hormone receptor NUR/77) (Nur77) (Orphan nuclear receptor HMR) (Orphan nuclear receptor TR3) (ST-59) (Testicular receptor 3)

Dilution

WB~~WB 1:500-2000, ELISA(peptide)1:5000-20000

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Nuclear Receptor NR4A1 (Phospho-Ser351) Antibody - Protein Information**Name** NR4A1**Synonyms** GFRP1, HMR, NAK1**Function**

Orphan nuclear receptor. Binds the NGFI-B response element (NBRE) 5'-AAAGGTCA-3' (PubMed:18690216, PubMed:8121493, PubMed:9315652). Binds 9-cis-retinoic acid outside of its ligand- binding (NR LBD) domain (PubMed:18690216). Participates in energy homeostasis by sequestering the kinase STK11 in the nucleus, thereby attenuating cytoplasmic AMPK activation (PubMed:22983157). Regulates the inflammatory response in macrophages by regulating metabolic adaptations during inflammation, including repressing the transcription of genes involved in the citric acid cycle (TCA) (By similarity). Inhibits NF-kappa-B signaling by binding to low-affinity NF-kappa-B binding sites, such as at the IL2 promoter (PubMed:18690216).

href="http://www.uniprot.org/citations/15466594" target="_blank">15466594). May act concomitantly with NR4A2 in regulating the expression of delayed-early genes during liver regeneration (By similarity). Plays a role in the vascular response to injury (By similarity).

Cellular Location

Nucleus. Cytoplasm, cytosol. Mitochondrion Note=Nuclear export to the cytosol is XPO1-mediated and positively regulated by IFI27 (PubMed:22427340). Translocation to the mitochondrion upon interaction with RXRA and upon the presence of 9-cis retinoic acid (PubMed:17761950).

Tissue Location


Fetal muscle and adult liver, brain and thyroid.

Nuclear Receptor NR4A1 (Phospho-Ser351) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Nuclear Receptor NR4A1 (Phospho-Ser351) Antibody - Images



Western blot analysis of various lysates, primary antibody was diluted at 1:1000, 4° over night, secondary antibody(cat# RS23920)was diluted at 1:10000, 37° 1hour.

Nuclear Receptor NR4A1 (Phospho-Ser351) Antibody - Background

Orphan nuclear receptor. May act concomitantly with NURR1 in regulating the expression of delayed-early genes during liver regeneration. Binds the NGFI-B response element (NBRE) 5'-AAAAGGTCA-3' (By similarity). May inhibit NF-kappa-B transactivation of IL2. Participates in energy homeostasis by sequestering the kinase STK11 in the nucleus, thereby attenuating cytoplasmic AMPK activation. Plays a role in the vascular response to injury (By similarity).