

14-3-3 ε Polyclonal Antibody

Catalog # AP68188

Specification

14-3-3 ε Polyclonal Antibody - Product Information

Application WB. IHC-P. IF **Primary Accession** P62258 Reactivity Human, Mouse, Rat Host Rabbit Clonality **Polyclonal**

14-3-3 ε Polyclonal Antibody - Additional Information

Gene ID 7531

Other Names

YWHAE; 14-3-3 protein epsilon; 14-3-3E

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence:

1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

IHC-P~~N/A IF~~1:50~200

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

14-3-3 ε Polyclonal Antibody - Protein Information

Name YWHAE

Function

Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways (PubMed:21189250). Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif (PubMed: 35343654). Binding generally results in the modulation of the activity of the binding partner (By similarity). Positively

regulates phosphorylated protein HSF1 nuclear export to the cytoplasm (PubMed: 12917326). Plays a positive role in the antiviral signaling pathway upstream of TBK1 via interaction with RIGI (PubMed:37555661). Mechanistically, directs RIGI redistribution from the cytosol to mitochondrial associated membranes where it mediates MAVS-dependent innate immune signaling during viral infection (PubMed:22607805).





Tel: 858.875.1900 Fax: 858.875.1999

Plays a role in proliferation inhibition and cell cycle arrest by exporting HNRNPC from the nucleus to the cytoplasm to be degraded by ubiquitination (PubMed:37599448).

Cellular Location


Nucleus. Cytoplasm Melanosome Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

14-3-3 ε Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

14-3-3 ε Polyclonal Antibody - Images







14-3-3 ε Polyclonal Antibody - Background

Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner (By similarity). Positively regulates phosphorylated protein HSF1 nuclear export to the cytoplasm (PubMed:12917326).