

ACADM Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6827c

Specification

ACADM Antibody (Center) - Product Information

Application IF, FC, IHC-P, WB,E

Primary Accession P11310
Other Accession O8HXY8

Reactivity Human, Mouse

Predicted Monkey
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 46588
Antigen Region 189-217

ACADM Antibody (Center) - Additional Information

Gene ID 34

Other Names

Medium-chain specific acyl-CoA dehydrogenase, mitochondrial, MCAD, ACADM

Target/Specificity

This ACADM antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 189-217 amino acids from the Central region of human ACADM.

Dilution

IF~~1:10~50 FC~~1:10~50 IHC-P~~1:50~100 WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ACADM Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ACADM Antibody (Center) - Protein Information





Name ACADM (HGNC:89)

Function Medium-chain specific acyl-CoA dehydrogenase is one of the acyl-CoA dehydrogenases that catalyze the first step of mitochondrial fatty acid beta-oxidation, an aerobic process breaking down fatty acids into acetyl-CoA and allowing the production of energy from fats (PubMed:1970566, PubMed:21237683, PubMed:2251268, PubMed:8823175). The first step of fatty acid beta-oxidation consists in the removal of one hydrogen from C-2 and C-3 of the straight-chain fatty acyl-CoA thioester, resulting in the formation of trans-2-enoyl-CoA (PubMed:2251268). Electron transfer flavoprotein (ETF) is the electron acceptor that transfers electrons to the main mitochondrial respiratory chain via ETF-ubiquinone oxidoreductase (ETF dehydrogenase) (PubMed:15159392, PubMed:25416781). Among the different mitochondrial acyl-CoA dehydrogenases, medium-chain specific acyl-CoA dehydrogenase acts specifically on acyl-CoAs with saturated 6 to 12 carbons long primary chains (PubMed:1970566, PubMed:21237683, PubMed:2251268, PubMed:8823175).


Cellular LocationMitochondrion matrix

ACADM Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ACADM Antibody (Center) - Images




Confocal immunofluorescent analysis of ACADM Antibody (Center)(Cat#AP6827c) with HepG2 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).






Western blot analysis of ACADM Antibody (Center) (Cat. #AP6827c) in mouse heart tissue lysates (35ug/lane). ACADM (arrow) was detected using the purified Pab.




Formalin-fixed and paraffin-embedded human colon carcinoma reacted with ACADM Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of widr cells using ACADM Antibody (Center)(bottom histogram) compared to a negative control cell (top histogram)FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ACADM Antibody (Center) - Background

ACADM is the medium-chain specific (C4 to C12 straight chain) acyl-Coenzyme A dehydrogenase. The homotetramer enzyme catalyzes the initial step of the mitochondrial fatty acid beta-oxidation





pathway.

ACADM Antibody (Center) - References

Ferreira, A.C., et.al., Genet. Mol. Res. 8 (2), 487-493 (2009)