

## **GNAS Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP6865c

## Specification

# **GNAS Antibody (Center) - Product Information**

WB,E
<u>Q5JWF2</u>
Human
Rabbit
Polyclonal
Rabbit IgG
111025
640-667

# **GNAS Antibody (Center) - Additional Information**

### Gene ID 2778

### **Other Names**

Guanine nucleotide-binding protein G(s) subunit alpha isoforms XLas, Adenylate cyclase-stimulating G alpha protein, Extra large alphas protein, XLalphas, GNAS, GNAS1 {ECO:0000303|PubMed:9707596}

#### Target/Specificity

This GNAS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 640-667 amino acids from the Central region of human GNAS.

Dilution WB~~1:4000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

GNAS Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

# **GNAS Antibody (Center) - Protein Information**

## Name GNAS

Synonyms GNAS1 {ECO:0000303|PubMed:9707596}



**Function** Guanine nucleotide-binding proteins (G proteins) function as transducers in numerous signaling pathways controlled by G protein- coupled receptors (GPCRs). Signaling involves the activation of adenylyl cyclases, resulting in increased levels of the signaling molecule cAMP. GNAS functions downstream of several GPCRs, including beta-adrenergic receptors. XLas isoforms interact with the same set of receptors as GNAS isoforms (By similarity).

### **Cellular Location**

Cell membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:Q63803}. Apical cell membrane

# **GNAS Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

## GNAS Antibody (Center) - Images



Anti-GNAS Antibody (Center) at 1:4000 dilution + Ramos whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 111 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

# **GNAS Antibody (Center) - Background**

Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems. The G(s) protein is involved in hormonal regulation of adenylate cyclase: it activates the cyclase in response to beta-adrenergic stimuli. XLas isoforms interact with the same set of receptors as Gnas isoforms.

## **GNAS Antibody (Center) - References**



Liu,C., et.al., Eur Arch Otorhinolaryngol (2009) Nishihara,E., et.al., Endocr. J. 56 (6), 791-798 (2009)