

C/EBP ζ Polyclonal Antibody
Catalog # AP68742**Specification**

C/EBP ζ Polyclonal Antibody - Product Information

Application	WB, IHC-P, IF
Primary Accession	Q03701
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

C/EBP ζ Polyclonal Antibody - Additional Information**Gene ID** 10153**Other Names**

CEBPZ; CBF2; CCAAT/enhancer-binding protein zeta; CCAAT-box-binding transcription factor; CBF; CCAAT-binding factor

DilutionWB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
IHC-P~~N/A
IF~~1:50~200**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

C/EBP ζ Polyclonal Antibody - Protein Information**Name** CEBPZ**Synonyms** CBF2**Function**

Stimulates transcription from the HSP70 promoter.

Cellular Location


Nucleus.

C/EBP ζ Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C/EBP ζ Polyclonal Antibody - Images



Western Blot analysis of various cells using C/EBP ζ Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100 (4°, overnight). High-pressure and temperature Tris-EDTA, pH 8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.

C/EBP ζ Polyclonal Antibody - Background

Stimulates transcription from the HSP70 promoter.