

Calmodulin Polyclonal Antibody

Catalog # AP68795

Specification

Calmodulin Polyclonal Antibody - Product Information

Application WB, IHC-P, IF

Primary Accession <u>P62158</u>

Reactivity Human, Mouse, Rat

Host Rabbit Clonality Polyclonal

Calmodulin Polyclonal Antibody - Additional Information

Other Names

CALM1; CALM; CAM1; CALM2; CAM2; CAMB; CALM3; CALML2; CAM3; CAMC; CAMIII;

Calmodulin; CaM

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence:

1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.

IHC-P~~N/A IF~~1:50~200

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Calmodulin Polyclonal Antibody - Protein Information


Calmodulin Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Calmodulin Polyclonal Antibody - Images







Western Blot analysis of various cells using Calmodulin Polyclonal Antibody diluted at 1□2000




Western Blot analysis of HuvEc cells using Calmodulin Polyclonal Antibody diluted at 1□2000



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted





at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.