

Casein Kinase Iε Polyclonal Antibody
Catalog # AP68834**Specification****Casein Kinase Iε Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IF
Primary Accession	P49674
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

Casein Kinase Iε Polyclonal Antibody - Additional Information**Gene ID** 1454**Other Names**

CSNK1E; Casein kinase I isoform epsilon; CKI-epsilon; CKIε

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

IHC-P~~N/A

IF~~1:50~200

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Casein Kinase Iε Polyclonal Antibody - Protein Information**Name** CSNK1E**Function**

Casein kinases are operationally defined by their preferential utilization of acidic proteins such as caseins as substrates (Probable). Participates in Wnt signaling (PubMed:12556519, PubMed:23413191). Phosphorylates DVL1 (PubMed:12556519). Phosphorylates DVL2 (PubMed:23413191). Phosphorylates NEDD9/HEF1 (By similarity). Central component of the circadian clock (PubMed:16790549). In balance with PP1, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation (PubMed:15917222, PubMed:16790549). Controls

PER1 and PER2 nuclear transport and degradation (By similarity). Inhibits cytokine-induced granulocytic differentiation (PubMed:15070676).

Cellular Location

Cytoplasm. Nucleus.


Tissue Location

Expressed in all tissues examined, including brain, heart, lung, liver, pancreas, kidney, placenta and skeletal muscle Expressed in monocytes and lymphocytes but not in granulocytes

Casein Kinase Iε Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Casein Kinase Iε Polyclonal Antibody - Images**Casein Kinase Iε Polyclonal Antibody - Background**

Casein kinases are operationally defined by their preferential utilization of acidic proteins such as caseins as substrates. Can phosphorylate a large number of proteins. Participates in Wnt signaling. Phosphorylates DVL1 and DVL2. Central component of the circadian clock. In balance with PP1, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation. Controls PER1 and PER2 nuclear transport and degradation. Inhibits cytokine-induced granulocytic differentiation.