

**CD1e Polyclonal Antibody** 

Catalog # AP68919

### Specification

## **CD1e Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<u>P15812</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Cionality	Polycional

### **CD1e Polyclonal Antibody - Additional Information**

Gene ID 913

**Other Names** CD1E; T-cell surface glycoprotein CD1e; membrane-associated; hCD1e; R2G1; CD antigen CD1e

**Dilution** WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.

**Format** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions** -20°C

# **CD1e Polyclonal Antibody - Protein Information**

Name CD1E

#### Function

T-cell surface glycoprotein CD1e, soluble binds diacetylated lipids, including phosphatidyl inositides and diacylated sulfoglycolipids, and is required for the presentation of glycolipid antigens on the cell surface. The membrane-associated form is not active.

#### **Cellular Location**

[T-cell surface glycoprotein CD1e, membrane- associated]: Golgi apparatus membrane; Single-pass type I membrane protein. Early endosome. Late endosome. Note=Predominantly localized in the trans-Golgi network in immature dendritic cells, and as a cleaved, soluble protein in the lysosome lumen of mature dendritic cells

#### **Tissue Location**

Expressed on cortical thymocytes, dendritic cells, Langerhans cells, on certain T-cell leukemias, and in various other tissues.




# **CD1e Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## **CD1e Polyclonal Antibody - Images**



# CD1e Polyclonal Antibody - Background

T-cell surface glycoprotein CD1e, soluble binds diacetylated lipids, including phosphatidyl inositides and diacylated sulfoglycolipids, and is required for the presentation of glycolipid antigens on the cell surface. The membrane- associated form is not active.